



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

PHYSICS 0625/23

Paper 2 Core

October/November 2010
1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units. Take the weight of 1 kg to be 10 N (i.e. acceleration of free fall = $10 \,\text{m/s}^2$).

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

1 Drops of water from a cracked gutter fall past the window of an IGCSE Physics student's room, as shown in Fig. 1.1.




Fig. 1.1

The student uses a digital stopwatch to find the time between one drop and the next.

To do this he

sets the stopwatch to zero,

then, starts the stopwatch as a drop comes into view at the top of the window,

then, stops the stopwatch 40 drops later.

The appearance of the stopwatch after 40 drops is shown in Fig. 1.2.



Fig. 1.2

(a) State the reading on the stopwatch.

reading =s [1]

(b) Calculate the time interval between one drop and the next.

(c)	time =
(•)	
(d)	Using Fig. 1.1, estimate the time for a drop to fall from the top of the upper window to the ground.
	time = s [3]
(e)	Fig. 1.1 shows that the drops get further apart as they get closer to the ground.
	Explain why this happens.
	[1]
	[Total: 8]

2 An unstretched spring of overall length 50.0 mm is hung from a support, as shown in Fig. 2.1.




Fig. 2.1

Different loads are placed on the spring and the extension is measured each time.

(a) On Fig. 2.1, mark clearly the extension caused by the load.

[1]

(b) The extensions for different loads are given in the table below.

load/N	extension/mm
0	0
1.0	10.0
2.0	20.5
3.0	31.0
4.0	41.5

(i) On Fig. 2.2, plot these values, using dots in small circles (⊙), and draw the best straight line for the points. [3]




Fig. 2.2

(ii) Complete the following sentence by inserting the appropriate word.

Within the limits of experimental accuracy, the load and the extension of the spring are to each other. [1]

(iii) A load of 2.5 N is hung on the spring.

1. What does the letter N stand for? [1]

2. Use the graph to estimate the overall length of the spring when 2.5 N is hanging from it.

length = mm [2]

[Total: 8]

3 (a) An aeroplane is flying horizontally at a steady speed in a straight line.

Fig. 3.1 shows three of the four forces acting on it.




Fig. 3.1

and are equal.	[1]
In order to fly horizontally in a straight line, there must be a fourth force acplane.	ting on the
Draw an arrow on Fig. 3.1 to represent this force.	[1]
	In order to fly horizontally in a straight line, there must be a fourth force ac plane.

(i) In order to fly horizontally at a steady speed, which two of the forces shown on the

- **(b)** The aeroplane in Fig. 3.1 flies an outward journey from Budapest (Hungary) to Palermo (Italy) in 2.75 hours. The distance is 2200 km.
 - (i) Calculate, in km/h, the average speed of the aeroplane.

average speed = km/h [3]

(ii) On the return journey from Palermo to Budapest, the journey time is shorter, even though the engine thrust is the same.

Suggest what might have caused the return journey to be shorter.

[41]

[Total: 6]

4 A simple pendulum starts with its bob at position X, shown in Fig. 4.1. The bob is pulled aside to Y and then released. It swings from Y to Z and back to Y.




Fig. 4.1

Write suitable words in the gaps in the following sentences. Ignore air resistance.

In order	to n	nove	the bob	from X to	Υ, .				has	to be	done	on	it and	lits
				energy	incr	eases	becau	se it	is	raised	furth	ner	from	the
ground.	As	it	moves	towards	Χ,	some	of	this	ene	ergy is	s co	nver	ted	into
			e	energy. Thro	ough	out the	swing	from	Y to	Z and	back	to Y	, the t	total
energy is				E	Enerç	gy is mo	easure	d in ur	nits ca	ılled				 [5]

[Total: 5]

5 (a) The list below contains terms that are used when dealing with heat and temperature.						heat and temperature.	
		ı	boiling point,	melting point,	internal energy,	thermal capacity	
		(i)	Which one of the	ese quantities will in	crease when an objec	ct is heated?	
							[1
		(ii)		ese determines the ut changing state?	temperature rise whe	en an object is given a quant	ity
							[1
	((iii)	Some liquid is he	eated until its tempe	erature stops rising.		
			Which one of the	ese quantities descr	ibes the temperature	at which this happens?	

(b) Fig. 5.1 shows an apparatus containing a brass rod. The brass rod is inside a tube, called a steam jacket, through which steam may be passed. The rod is fixed at the right-hand end, but free to move at the left-hand end. The dial micrometer indicates any movement of the left hand end.




Fig. 5.1

Steam is now passed through the steam jacket. In the boxes below, write down what will happen to the readings on the thermometer and the dial micrometer, and why.

	what will happen	why
reading on thermometer		
reading on dial micrometer		

[4]

6 (a) Fig. 6.1 shows a ray of light AB striking a plane mirror at an angle of incidence of 40°.




Fig. 6.1

State the value of the angle of reflection of the ray[1]

(b) In Fig. 6.2, the mirror has been rotated 10° from its position in Fig. 6.1. AB has remained unchanged.




Fig. 6.2

(i) On Fig. 6.2, use a straight edge to draw the reflected ray.

(ii) State the value of the angle between the reflected ray and the line BC.

(c) An object of height 2cm is placed 5cm in front of a plane mirror.

(i) State the height of the image formed by the mirror. [1]

(ii) Find the distance between the object and the image.

distance = cm [2]

[Total: 8]

7 A narrow beam of white light enters a glass prism and is split into the colours of the visible spectrum, as shown (not to scale) in Fig. 7.1.




Fig. 7.1

(a)	Wha	What name do we give to								
	(i)	the bending of the light as it enters the prism,								
	(ii)	the different amounts of bending that give rise to the spectrum?								
		[2								
(b)	The	lines leaving the prism represent rays of the seven main colours of the visible spectrum								
	In th	ne answer spaces provided on Fig. 7.1, write								

(c) The visible spectrum is part of the electromagnetic spectrum.

'yellow' in the space alongside the yellow ray.

'red' in the space alongside the red ray,

(ii)

[Total: 6]

[2]

8 A stretched string is vibrating between two fixed ends. Fig. 8.1 shows how the string is vibrating.




Fig. 8.1

		1.9.011	
(a)	Stat	te the name of	
	(i)	distance a,	[1]
	(ii)	distance b.	[1]
(b)	The	string is causing a sound to be transmitted through the air.	
	(i)	Describe how the string causes the sound.	
			[2]
	(ii)	State what happens to the sound as the distance a decreases.	
			[1]
		т]	otal: 5]

9	(a)	(i)	In the space below, draw a diagram of the circuit that you would use to determine the
			resistance of a coil of wire using a voltmeter and an ammeter.

Use conventional symbols and label the coil clearly.

[3]

(ii) State the equation you would use to calculate the resistance of the coil.

[1]

(iii) State two properties of the wire on which the resistance of the coil depends.

1.

2......[2]

(b) In Fig. 9.1, AB is a 2.0 m length of uniform resistance wire, connected into a circuit.

Ignore the resistance of the battery.




Fig. 9.1

The current	I	is	1	.5	A.
-------------	---	----	---	----	----

Calculate the resistance per metre of the resistance wire.

resistance per metre = Ω/m [4]

[Total: 10]

10 (a) The apparatus in Fig. 10.1 is set up in a laboratory. The metal wheels are rolled along the rails from the left-hand end to the right-hand end.




Fig. 10.1

(i)	Describe what is seen happening to the pointer on the sensitive centre-zero millivoltmeter.
	[2]
(ii)	Explain why this happens.
	[3]
(iii)	The metal wheels are now rolled back to the left-hand end again.
	Describe what now happens to the millivoltmeter pointer.
	[1]

(b) Fig. 10.2 shows a magnet suspended above a coil of wire.




Fig. 10.2

The end of the magnet is pushed into the coil and released, so that it bounces repeatedly in and out of the coil.

Describe what is seen on the sensitive centre-zero millivoltmeter.	
	[1]
	[Total: 7]

(a) In the space below, draw the circuit symbol for a fuse.

[1]

[Total: 4]

12	(a)	A β-particle may be represented by the symbol $_{-1}^{0}$ e.			
		(i)	What does the e indicate about a β-particle?		
		(ii)	What does the 0 indicate about a β-particle?		
		(iii)	What does the -1 indicate about a β-particle?		
				Г 4	

(b) The nuclide $^{250}_{\ \, 97}Bk$ decays by emitting a $\beta\text{-particle}.$

Complete the nuclear equation for this decay by writing appropriate numbers in the boxes.

$$^{250}_{97}Bk \longrightarrow Cf + ^{0}_{-1}e$$

[Total: 6]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.