

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

**0620/33**

Paper 3 (Extended)

**October/November 2015**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

1 (a) Describe a chemical test which shows the presence of water.

test .....

colour change if water is present .....

..... [3]

(b) How could you show that a sample of water is pure?

..... [1]

(c) Describe how water is treated before it is supplied to homes and industry.

.....

..... [2]

(d) State **two** industrial uses of water.

.....

..... [2]

[Total: 8]

2 Choose from the following list of gases. A gas may be chosen once, more than once or not at all.

**sulfur dioxide**

**hydrogen**

**methane**

**carbon monoxide**

**argon**

**ethene**

**butane**

(a) It is used to bleach wood pulp. .... [1]

(b) When burned in oxygen, the only product is water. .... [1]

(c) It can polymerise. .... [1]

(d) It is used to provide an inert atmosphere for welding. .... [1]

(e) When reacted with oxygen, the only product is carbon dioxide. .... [1]

(f) It is produced by the decay of vegetation in the absence of oxygen. .... [1]

[Total: 6]

3 Lithium bromide is an ionic compound. It can be electrolysed when it is molten or in aqueous solution. It cannot be electrolysed as a solid.

(a) Solid lithium bromide is a poor conductor of electricity. The ions cannot move to the electrodes, they are held in an ionic lattice by strong forces.

(i) Describe the motion of the ions in the solid state.

..... [1]

(ii) Define the term *ionic bonding*.

.....


..... [2]

(iii) What is meant by the term *ionic lattice*?

.....

..... [2]

(b) The diagram shows the electrolysis of molten lithium bromide.



(i) Mark on the diagram the direction of the electron flow. [1]

(ii) Write an ionic equation for the reaction at the negative electrode (cathode).

..... [1]

(iii) Write an ionic equation for the reaction at the positive electrode (anode).

..... [2]

(iv) Which ion is oxidised? Explain your answer.

.....

..... [2]

- (c) When aqueous lithium bromide is electrolysed, a colourless gas is formed at the negative electrode and the solution becomes alkaline.

Explain these observations and include an equation in your explanation.

.....

.....

.....

..... [3]

[Total: 14]

- 4 Two homologous series of hydrocarbons are the alkanes and the alkenes.

- (a) (i) One general characteristic of a homologous series is that the physical properties vary in a predictable way.

State **three** other general characteristics of a homologous series.

.....

.....

..... [3]

- (ii) How can the molecular formula of a hydrocarbon show whether it is an alkane or an alkene?

.....

..... [2]

- (iii) How do alkanes and alkenes differ in their molecular structures?

.....

..... [2]

**(b)** Cracking is the thermal decomposition of alkanes into smaller hydrocarbons and possibly hydrogen.

**(i)** State **two** conditions required for the cracking of an alkane.


..... [2]

**(ii)** One type of cracking produces an alkane and an alkene.

Complete an equation for the cracking of heptane into an alkane and an alkene.



**(iii)** Complete an equation for the cracking of heptane into hydrogen and two other products.



**(iv)** Suggest **one** reason why cracking is important.

..... [1]

**(c)** Hydrocarbons burn in excess oxygen to form carbon dioxide and water. 20 cm<sup>3</sup> of a gaseous hydrocarbon burned in an excess of oxygen, 200 cm<sup>3</sup>. After cooling, the volume of the residual gas at r.t.p. was 150 cm<sup>3</sup>, 50 cm<sup>3</sup> of which was oxygen.

**(i)** Determine the volume of the oxygen used.

..... [1]

**(ii)** Determine the volume of the carbon dioxide formed.

..... [1]

**(iii)** The hydrocarbon was an alkane.

Determine the formula of the hydrocarbon.

[1]

[Total: 15]

5 Sulfuric acid is a strong acid. In aqueous solution, it ionises as shown below.



(a) (i) What is meant by the term *acid*?

..... [1]

(ii) Sulfurous acid,  $\text{H}_2\text{SO}_3$ , is a weak acid.

State the difference between a weak acid and a strong acid.

.....  
 ..... [2]

(b) Sulfurous acid forms salts called sulfites, which contain the ion  $\text{SO}_3^{2-}$ .

When barium nitrate solution is added to aqueous sulfurous acid, a white precipitate, **A**, forms.

Bromine water changes from brown to colourless when added to aqueous sulfurous acid.

Bromine oxidises sulfurous acid. When this solution is tested with acidified barium nitrate solution, a different white precipitate, **B**, is formed.

(i) Identify the white precipitate, **A**.

..... [1]

(ii) Identify the white precipitate, **B**.

..... [1]

(iii) Write an ionic equation for the reduction of the bromine molecule.

..... [1]

(iv) Name the product formed by the oxidation of sulfurous acid.

..... [1]

(c) Complete the following word equations.

(i) magnesium hydroxide + dilute sulfuric acid

..... [1]

(ii) zinc + dilute sulfuric acid

..... [1]

(iii) copper carbonate + dilute sulfuric acid

..... [1]

(d) Write equations for the reaction of dilute sulfuric acid with each of the following.

(i) ammonia

..... [2]

(ii) sodium hydroxide

..... [2]

(iii) iron

..... [2]

[Total: 16]

6 A reactivity series of metals is given below.

	metal name	symbol
most reactive ↓ least reactive	sodium	Na
	lithium	Li
	magnesium	Mg
	zinc	Zn
	manganese	Mn
	iron	Fe
	copper	Cu
	rhodium	Rh

(a) Which **two** metals will react most vigorously with cold water?

..... [1]

(b) Which **two** metals will not react with dilute hydrochloric acid?

..... [1]

(c) Deduce the formula of iron(III) sulfate.

..... [1]

(d) What is the formula of a magnesium ion?

..... [1]

(e) Describe a test-tube experiment which will show that manganese is more reactive than copper.

.....  
 .....  
 ..... [3]



(f) Manganese is a typical transition metal.

Predict **three** physical and **two** chemical properties of this metal.

physical properties

.....  
.....  
.....


chemical properties

.....  
.....


[5]

[Total: 12]

- 7 Two salts can be made from potassium hydroxide and sulfuric acid. They are potassium sulfate,  $K_2SO_4$ , and the acid salt potassium hydrogen sulfate,  $KHSO_4$ . They are both made by titration.



- (a) 25.0 cm<sup>3</sup> of potassium hydroxide, concentration 2.53 mol/dm<sup>3</sup>, was neutralised by 28.2 cm<sup>3</sup> of dilute sulfuric acid.



Calculate the concentration of the sulfuric acid.

number of moles of KOH used = .....

number of moles of  $H_2SO_4$  needed to neutralise the KOH = .....

concentration of dilute sulfuric acid = ..... mol/dm<sup>3</sup>


[3]

- (b) In the conical flask there is a neutral solution of potassium sulfate which still contains the indicator used in the titration.

- (i) Describe how you could obtain a solution of potassium sulfate without the indicator.

.....  
 ..... [2]

- (ii) Potassium hydrogen sulfate can be made by the following reaction.



Suggest how you could make a solution of potassium hydrogen sulfate without using an indicator.

.....  
 .....  
 ..... [2]

(c) Describe a test which would distinguish between aqueous solutions of potassium sulfate and sulfuric acid.

test .....

result .....

[2]

[Total: 9]

## DATA SHEET The Periodic Table of the Elements

Group																				
I	II											III	IV	V	VI	VII	0			
												1 <b>H</b> Hydrogen 1								4 <b>He</b> Helium 2
7 <b>Li</b> Lithium 3	9 <b>Be</b> Beryllium 4											11 <b>B</b> Boron 5	12 <b>C</b> Carbon 6	14 <b>N</b> Nitrogen 7	16 <b>O</b> Oxygen 8	19 <b>F</b> Fluorine 9	20 <b>Ne</b> Neon 10			
23 <b>Na</b> Sodium 11	24 <b>Mg</b> Magnesium 12											27 <b>Al</b> Aluminium 13	28 <b>Si</b> Silicon 14	31 <b>P</b> Phosphorus 15	32 <b>S</b> Sulfur 16	35.5 <b>Cl</b> Chlorine 17	40 <b>Ar</b> Argon 18			
39 <b>K</b> Potassium 19	40 <b>Ca</b> Calcium 20	45 <b>Sc</b> Scandium 21	48 <b>Ti</b> Titanium 22	51 <b>V</b> Vanadium 23	52 <b>Cr</b> Chromium 24	55 <b>Mn</b> Manganese 25	56 <b>Fe</b> Iron 26	59 <b>Co</b> Cobalt 27	59 <b>Ni</b> Nickel 28	64 <b>Cu</b> Copper 29	65 <b>Zn</b> Zinc 30	70 <b>Ga</b> Gallium 31	73 <b>Ge</b> Germanium 32	75 <b>As</b> Arsenic 33	79 <b>Se</b> Selenium 34	80 <b>Br</b> Bromine 35	84 <b>Kr</b> Krypton 36			
85 <b>Rb</b> Rubidium 37	88 <b>Sr</b> Strontium 38	89 <b>Y</b> Yttrium 39	91 <b>Zr</b> Zirconium 40	93 <b>Nb</b> Niobium 41	96 <b>Mo</b> Molybdenum 42	96 <b>Tc</b> Technetium 43	101 <b>Ru</b> Ruthenium 44	103 <b>Rh</b> Rhodium 45	106 <b>Pd</b> Palladium 46	108 <b>Ag</b> Silver 47	112 <b>Cd</b> Cadmium 48	115 <b>In</b> Indium 49	119 <b>Sn</b> Tin 50	122 <b>Sb</b> Antimony 51	128 <b>Te</b> Tellurium 52	127 <b>I</b> Iodine 53	131 <b>Xe</b> Xenon 54			
133 <b>Cs</b> Caesium 55	137 <b>Ba</b> Barium 56	139 <b>La</b> Lanthanum 57 *	178 <b>Hf</b> Hafnium 72	181 <b>Ta</b> Tantalum 73	184 <b>W</b> Tungsten 74	186 <b>Re</b> Rhenium 75	190 <b>Os</b> Osmium 76	192 <b>Ir</b> Iridium 77	195 <b>Pt</b> Platinum 78	197 <b>Au</b> Gold 79	201 <b>Hg</b> Mercury 80	204 <b>Tl</b> Thallium 81	207 <b>Pb</b> Lead 82	209 <b>Bi</b> Bismuth 83	209 <b>Po</b> Polonium 84	209 <b>At</b> Astatine 85	209 <b>Rn</b> Radon 86			
87 <b>Fr</b> Francium	226 <b>Ra</b> Radium 88	227 <b>Ac</b> Actinium 89 †																		
*58-71 Lanthanoid series			140 <b>Ce</b> Cerium 58	141 <b>Pr</b> Praseodymium 59	144 <b>Nd</b> Neodymium 60	147 <b>Pm</b> Promethium 61	150 <b>Sm</b> Samarium 62	152 <b>Eu</b> Europium 63	157 <b>Gd</b> Gadolinium 64	159 <b>Tb</b> Terbium 65	162 <b>Dy</b> Dysprosium 66	165 <b>Ho</b> Holmium 67	167 <b>Er</b> Erbium 68	169 <b>Tm</b> Thulium 69	173 <b>Yb</b> Ytterbium 70	175 <b>Lu</b> Lutetium 71				
†90-103 Actinoid series			232 <b>Th</b> Thorium 90	234 <b>Pa</b> Protactinium 91	238 <b>U</b> Uranium 92	237 <b>Np</b> Neptunium 93	244 <b>Pu</b> Plutonium 94	247 <b>Am</b> Americium 95	251 <b>Cm</b> Curium 96	252 <b>Bk</b> Berkelium 97	257 <b>Cf</b> Californium 98	261 <b>Es</b> Einsteinium 99	267 <b>Fm</b> Fermium 100	271 <b>Md</b> Mendelevium 101	277 <b>No</b> Nobelium 102	289 <b>Lr</b> Lawrencium 103				

Key

a
<b>X</b>
b

a = relative atomic mass  
X = atomic symbol  
b = proton (atomic) number

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).