



### **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/32
Paper 3 (Extend	ded)	F	ebruary/March 2015
			1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1		ches the description.
	(a)	an element which is gaseous at room temperature and pressure
		[1]
	(b)	an element that is added to water to kill bacteria
		[1]
	(c)	an element that forms a basic oxide of the type XO
		[1]
	(d)	an element used as an inert atmosphere in lamps
	(0)	an element that forms an amphatoria axide
	(e)	an element that forms an amphoteric oxide[1]
	(f)	an element that reacts vigorously with cold water to produce hydrogen
	(-)	[1]
		[Total: 6]
2	(a)	Define the term <i>isotope</i> .
		[2]
	(ls)	The table vives information about four portions A. D. C. and D.
	(D)	The table gives information about four particles, <b>A</b> , <b>B</b> , <b>C</b> and <b>D</b> .
		Complete the table. The first line has been done for you.

particle	number of protons	number of electrons	number of neutrons	nucleon number	symbol or formula
Α	6	6	6	12	С
В	11	10	12		
С	8		8		O <sup>2-</sup>
D		10		28	Al <sup>3+</sup>

[7]

[Total: 9]

Ammonia is manufactured by the Haber process. Nitrogen and hydrogen are passed over a catalyst at a temperature of 450 $^{\circ}$ C and a pressure of 200 atmospheres.

The equation for the reaction is as follows.

$$N_2 + 3H_2 \rightleftharpoons 2NH_3$$

The forward reaction is exothermic.

(a)	Sta	te <b>one</b> use of ammonia. [1
(b)	Wh	at is the meaning of the symbol <del>←</del> ?
(c)		at are the sources of nitrogen and hydrogen used in the Haber process?
	hyd	rogen[2]
(d)	Nar	me the catalyst in the Haber process.
(e)	(i)	If a temperature higher than 450 °C was used in the Haber process, what would happen to the <b>rate</b> of the reaction? Give a reason for your answer.
	(ii)	If a temperature higher than 450 °C was used in the Haber process, what would happen to the <b>yield</b> of ammonia? Give a reason for your answer.

(f)	(i)	If a pressure higher than 200 atmospheres was used in the Haber process, what would happen to the <b>yield</b> of ammonia? Give a reason for your answer.	d
		[2	
	(ii)	Explain why the rate of reaction would be faster if the pressure was greater tha 200 atmospheres.	n
	(iii)	Suggest <b>one</b> reason why a pressure higher than 200 atmospheres is not used in the Haber process.	
		[1	
(g)		w a dot-and-cross diagram to show the arrangement of the outer (valency) electrons in one lecule of ammonia.	1)
		[2	<u>?]</u>
(h)	Am	monia acts as a base when it reacts with sulfuric acid.	
	(i)	What is a base?	
	(ii)	Write a balanced equation for the reaction between ammonia and sulfuric acid.	
		[2	-]

(a)	A) A compound <b>X</b> contains 82.76% of carbon by mass and 17.24% of hydrogen by mass.		
	(i)	Calculate the empirical formula of compound <b>X</b> .	
	(ii)	Compound <b>X</b> has a relative molecular mass of 58.	[2]
	(11)		
		Deduce the molecular formula of compound <b>X</b> .	[2]
(b)	Alk	enes are unsaturated hydrocarbons.	
	(i)	State the general formula of alkenes.	[41
	(ii)	State the empirical formula of alkenes.	ניו
			[1]
(c)		at is meant by the term unsaturated hydrocarbon?	
	hya	rocarbon	
			 [2]

(d)	Des	escribe a test that would distinguish between saturated and unsaturated hydrocarbons.		
	rea	gent		
	obs	ervation (saturated hydrocarbon)		
	obs	ervation (unsaturated hydrocarbon)[3]		
(e)	Add	dition polymers can be made from alkenes. The diagram shows part of an addition polymer.		
		$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		
	(i)	Draw a circle on the diagram to show one repeat unit in this polymer. [1]		
	(ii)	Give the structure and the name of the monomer used to make this polymer.		
		structure		

name [2	2]
---------	----


(iii) Give the structure of an isomer of the alkene in (e)(ii).

[1]

[Total: 15]


**5** Aluminium and iron are extracted from their ores by different methods.

Aluminium is extracted from its purified oxide ore by electrolysis.



(HCţ	aluminium
(a)	What is the name of the ore of aluminium which consists mainly of aluminium oxide?
	[1]
(b)	The electrodes are both made of the same substance.
	Name this substance.
	[1]
(c)	Aluminium oxide is dissolved in molten cryolite before it is electrolysed.
	Give <b>two</b> reasons why aluminium oxide dissolved in molten cryolite is electrolysed rather than molten aluminium oxide alone.
	[2]
(d)	Write the <b>ionic</b> equations for the reactions at the electrodes in this electrolysis.
	anode (positive electrode)
	cathode (negative electrode)

(e) Iron is extracted from its oxide ore by reduction using carbon in a blast furnace.



	(i)	Place the elements aluminium, carbon and iron in order of reactivity with the <b>least</b> reactive element first.	ive
	(ii)	Use your answer to (e)(i) to explain why iron is extracted by reduction using carbon by aluminium is not.	but
(f)	Wh	at is the name of the ore of iron which consists mainly of iron(III) oxide?	[1]
(g)	Wri	te balanced equations for the reactions occurring in the blast furnace which involve the complete combustion of coke (carbon),	
	(1)	the complete combustion of coke (carbon),	[1]
	(ii)	the production of carbon monoxide from carbon dioxide,	[1]
	(iii)	the reduction of iron(III) oxide,	[11
	(iv)	the formation of slag.	ניו
			[1]

**6** A student is told to produce the maximum amount of copper from a mixture of copper and copper(II) carbonate.

The student adds the mixture to an excess of dilute sulfuric acid in a beaker and stirs the mixture with a glass rod. The copper(II) carbonate reacts with the sulfuric acid, forming a solution of copper(II) sulfate but the copper does not react with the sulfuric acid.

#### The student then

Torrioted the armedoted dopper morn the mixture,	•	removes	the unreacted	copper from	the mixture,
--	---	---------	---------------	-------------	--------------

<ul> <li>CO</li> </ul>	nverts the	solution c	of copper(	(II) sulfate	e into coppe	er by a	series of	reactions.
------------------------	------------	------------	------------	--------------	--------------	---------	-----------	------------

(a)		scribe <b>two</b> things that the student would observe when the mixture is added to the dilute furic acid.
		[2]
(b)		scribe how the student can produce pure dry copper from the mixture of copper and $\mbox{\rm sper}(\Pi)$ sulfate solution.
		[3]
(c)		e student then adds sodium hydroxide solution to the copper(II) sulfate solution to produce $per(\mathrm{II})$ hydroxide.
	(i)	Describe what the student would observe.
		[1]
	(ii)	Write an <b>ionic</b> equation for this reaction.
		[1]
(d)		er separating the copper(II) hydroxide from the mixture, the copper(II) hydroxide is heated engly. The copper(II) hydroxide decomposes into copper(II) oxide and steam.
	(i)	Write an equation for the decomposition of copper(II) hydroxide. Include state symbols.
		[2]
	(ii)	Name a non-metallic element that can be used to convert copper(II) oxide into copper.
		[1]

[Total: 10]

 $Ethanol\ is\ manufactured\ from\ glucose,\ C_6H_{12}O_6,\ by\ fermentation\ according\ to\ the\ following\ equation.$ 

		$C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$	
(a)	Sta	te the conditions required for this reaction.	
			[2]
(b)	In a	an experiment, 30.0g of glucose was fermented.	
	(i)	Calculate the number of moles of glucose in 30.0 g.	
		mol	[2]
(	(ii)	Calculate the maximum mass of ethanol that could be obtained from 30.0 g of glucose.	
		g	[2]
(	iii)	Calculate the volume of carbon dioxide at room temperature and pressure that can l	
		obtained from 30.0 g of glucose.	
		dm³	[1]
(c)	Eth	anol can also be manufactured from ethene.	
	(i)	Name the raw material which is the source of ethene.	
			[1]
(	(ii)	Write a balanced equation for the manufacture of ethanol from ethene.	
			[1]
		[Total:	9]

## **BLANK PAGE**

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

# DATA SHEET The Periodic Table of the Elements

Group																		
I	II											III	IV	V	VI	VII	0	
		1 H Hydrogen										4 He Helium 2						
7	9							-				11	12	14	16	19	20	
Li Lithium	Be Beryllium											B Boron	C Carbon 6	N Nitrogen	Oxygen 8	F Fluorine 9	Ne Neon	
23 Na Sodium	24 Mg Magnesium 12											27 Al Aluminium 13	28 Si Silicon	31 P Phosphorus 15	32 <b>S</b> Sulfur	35.5 <b>C1</b> Chlorine	40 Ar Argon	
39 <b>K</b> Potassium 19	40 Ca Calcium	45 Sc Scandium 21	48 <b>Ti</b> Titanium 22	51 <b>V</b> Vanadium 23	52 Cr Chromium 24	55 Mn Manganese 25	56 <b>Fe</b> Iron	59 Co Cobalt 27	59 <b>Ni</b> Nickel	64 Cu Copper 29	65 <b>Zn</b> Zinc	70 <b>Ga</b> Gallium	73 <b>Ge</b> Germanium 32	75 <b>As</b> Arsenic	79 Se Selenium 34	80 <b>Br</b> Bromine 35	84 <b>Kr</b> Krypton 36	
85 <b>Rb</b> Rubidium 37	88 Sr Strontium 38	89 <b>Y</b> Yttrium	91 <b>Zr</b> Zirconium 40	93 <b>Nb</b> Niobium	96 Mo Molybdenum 42	Tc Technetium 43	101 <b>Ru</b> Ruthenium 44	103 <b>Rh</b> Rhodium 45	106 Pd Palladium 46	108 <b>Ag</b> Silver	112 Cd Cadmium 48	115 <b>In</b> Indium 49	119 <b>Sn</b> Tin	122 <b>Sb</b> Antimony 51	128 <b>Te</b> Tellurium 52	127 <b>I</b> lodine 53	131 <b>Xe</b> Xenon 54	
133 Cs Caesium 55	137 <b>Ba</b> Barium	139 <b>La</b> Lanthanum  57 *	178 <b>Hf</b> Hafnium  72	181 <b>Ta</b> Tantalum 73	184 W Tungsten 74	186 <b>Re</b> Rhenium 75	190 Os Osmium 76	192 Ir Iridium	195 Pt Platinum 78	197 <b>Au</b> Gold	201 <b>Hg</b> Mercury	204 <b>T <i>l</i></b> Thallium 81	207 <b>Pb</b> Lead	209 <b>Bi</b> Bismuth	Po Polonium 84	At Astatine 85	Rn Radon 86	
Fr Francium 87	226 <b>Ra</b> Radium	227 Ac Actinium 89 †			,													
*58-71 Lanthanoid series †90-103 Actinoid series			140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	Pm Promethium 61	150 Sm Samarium 62	152 <b>Eu</b> Europium 63	157 <b>Gd</b> Gadolinium 64	159 <b>Tb</b> Terbium 65	162 <b>Dy</b> Dysprosium 66	165 <b>Ho</b> Holmium 67	167 <b>Er</b> Erbium 68	169 <b>Tm</b> Thulium 69	173 <b>Yb</b> Ytterbium 70	175 Lu Lutetium 71		
Key X			232 <b>Th</b> Thorium 90	Pa Protactinium 91	238 <b>U</b> Uranium 92	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	<b>Bk</b> Berkelium  97	Cf Californium 98	<b>Es</b> Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103		

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).