Location Entry Codes



As part of CIE's continual commitment to maintaining best practice in assessment, CIE has begun to use different variants of some question papers for our most popular assessments with extremely large and widespread candidature, The question papers are closely related and the relationships between them have been thoroughly established using our assessment expertise. All versions of the paper give assessment of equal standard.

The content assessed by the examination papers and the type of questions are unchanged.

This change means that for this component there are now two variant Question Papers, Mark Schemes and Principal Examiner's Reports where previously there was only one. For any individual country, it is intended that only one variant is used. This document contains both variants which will give all Centres access to even more past examination material than is usually the case.

The diagram shows the relationship between the Question Papers, Mark Schemes and Principal Examiner's Reports.

Question Paper

Introduction First variant Question Paper Second variant Question Paper

Mark Scheme

| Introduction |
|----------------------------|
| First variant Mark Scheme |
| |
| Second variant Mark Scheme |

Principal Examiner's Report

| Introduction |
|--|
| First variant Principal Examiner's Report |
| Second variant Principal Examiner's Report |

Who can I contact for further information on these changes?

Please direct any questions about this to CIE's Customer Services team at: international@cie.org.uk

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the October/November 2008 question paper

0620 CHEMISTRY

0620/31

Paper 31 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2008 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



| Page 2 | Mark Scheme | Syllabus | Paper |
|--------|-------------------------------|----------|-------|
| | IGCSE – October/November 2008 | 0620 | 31 |

| 1 | | | us paper blue se fumes/smoke with HC $l\left(\mathrm{g}\right)$ or (aq) | [1] | | |
|---|--------|------------------------|--|---------------------------------|--|--|
| | chlo | orine | | [1] | | |
| | | • | th a lighted splint or burn with a pop or goes pop and extinguishes flame owing splint | [1] | | |
| | oxygen | | | | | |
| | | | dioxide T correct formulae | [1] | | |
| | | | | [Total: 5] | | |
| 2 | (a) | corı | a : 1N correct ratio rect charges around N | [1] [1] [1] | | |
| | | if co igno if th | o symbols then must have correct key ovalent only mark 1 ore electrons around sodium e response includes both a correct and an incorrect answer not select correct one, mark = [0] | | | |
| | (b) | (i) | positive ions or cations NOT atoms or cores or nuclei layers or lattice or regular pattern delocalised or free or mobile electrons or sea OR positive ions or cations NOT atoms or cores or nuclei attraction between ions and electrons delocalised or free or mobile electrons or sea | [1] [1] [1] [1] [1] | | |
| | | | the attraction/electrostatic bonding must be between ions and delocalised electrons, between cations and anions does not score ACCEPT bond if qualified - electrostatic bond, etc. if molecular or molecules then cannot score cation mark | [.1 | | |
| | | (ii) | delocalised/free/mobile electrons or electrons can move | [1] | | |
| | | | layers or ions or atoms or particles NB more flexible than 2(b)(i) can <u>slip</u> or move past each other or bonding non-directional | [1] [1] | | |

| Page 3 | Mark Scheme | Syllabus | Paper |
|---------|---------------------------------|----------|-------|
| | IGCSE – October/November 2008 | 0620 | 31 |
| (c) (i) | tetrahedral | | [′ |
| | 1Si: 4O bonded/surrounded, etc. | | [|

NOT molecules of oxygen, etc.

NOT intermolecular forces

ONLY tetrahedral can score for either of the above

Despite what the question states, ACCEPT a clear accurate diagram which shows the above three points.

[1]

[1]

(ii) hard

high mp or bp

10:2 Si

colourless (NOT clear) or shiny or translucent

non/poor conductor (of electricity)

brittle

insoluble

any TWO

[2]

NOT crystalline or strong

[Total: 14]

- 3 (a) (i) water or moisture ACCEPT salty water [1] air or oxygen [1]
 - (ii) galvanising or coat with zinc

tin plate

chromium plate

nickel plate

cobalt plate

copper plate

cover with aluminium

anodic protection or sacrificial protection

cathodic protection

cover with plastic

alloying (ignore any named metal)

any TWO [2]

NOT just plate or electroplate need electroplate with suitable metal

NOT oil

ACCEPT both galvanising and sacrificial protection

- (b) (i) hydrogen or carbon or carbon monoxide or methane or more reactive metal NOT Group I
 - (ii) any correct equation [2]

only error not balanced [1]

| | Page 4 | | · | Mark Scheme | Syllabus | Paper |
|---|--------|-------|--|---|----------|-------------|
| | | | | IGCSE – October/November 2008 | 0620 | 31 |
| | (c) | (i) | 196 | | | [1] |
| | | (ii) | 112/ = 57 mark ONL othe | [1] [1] | | |
| | (d) | (i) | form | ns carbon dioxide/carbon monoxide (which escapes) |) | [1] |
| | | (ii) | | ns silicon(IV) oxide or silicon oxide or silica | | [1] |
| | | | to fo igno | CaO reacts with SiO ₂ orm slag or calcium silicate ore an incorrect formula if a correct name "slag" give r Si + O ₂ + CaO form slag, this gains mark for slag o | | [1] |
| | | | | | | [Total: 13] |
| 4 | (a) | (i) | | $_{5}$ COOH or C_{6} H $_{5}$ CO $_{2}$ H Γ C_{7} H $_{6}$ O $_{2}$ / C_{6} H $_{6}$ COO | | [1] |
| | | (ii) | corre | um hydroxide + benzoic acid = sodium benzoate + vect spelling needed NOT benzenoate CEPT correct symbol equation | water | [1] |
| | | (iii) | | um carbonate or oxide or hydrogencarbonate TWO Γ Na | | [2] |
| | (b) | (i) | 7.7% | % | | [1] |
| | | (ii) | | any number: equal number ratio example 1:1 or 6:6 | | [2] |
| | | (iii) | mole | pirical formula is CH ecular formula is C_6H_6 e.c.f., award of marks not dependent on (ii) | | [1] [1] |
| | (c) | (i) | C ₆ H ₈ | ₈ O ₆ | | [1] |
| | | (ii) | alco NOT | oon – carbon double bond or alkene hol or hydroxyl or hydroxy I hydroxide roxide and alcohol = 0 | | [1] [1] |
| | | | | | | [Total: 12] |

| Page 5 | Mark Scheme | Syllabus | Paper |
|--------|-------------------------------|----------|-------|
| | IGCSE – October/November 2008 | 0620 | 31 |

- 5 (a) (i) $2H^+ + 2e \rightarrow H_2$ [1]
 - (ii) $2Cl^- 2e \rightarrow Cl_2$ or $2Cl^- \rightarrow Cl_2 + 2e$ [1]
 - (iii) Na⁺ and OH⁻ are left
 OR C*l*⁻ removed OH⁻ left

 NB ions by name or formula essential

NB ions by name **or** formula essential **NOT** any reaction of Na **or** Na⁺ **NOT** Na⁺ and OH⁻ combine

- (b) (i) sterilise/disinfect water or kill microbes/germs bacteria, etc.
 NOT just to make it safe to drink or purify it or clean it treat above as neutral they do not negate a correct response
 - (ii) ammonia **or** methanol **or** hydrogen chloride **or** margarine [1] **NOT** nylon
 - (iii) fat or lipid or triester or named fat or glyceryl stearate

 or vegetable oil

 heat

 [1]

[Total: 7]

6 (a) (i)

| aqueous solution | tin Sn | manganese Mn | silver Ag | zinc Zn |
|-----------------------|-----------|-----------------|--------------|------------|
| tin(II) nitrate | | R | NR | R |
| manganese(II) nitrate | NR | | NR | NR |
| silver(I) nitrate | R | R | | R |
| zinc nitrate | NR | R | NR | |

- [1] for each row [3] ignore anything written in blank space
- (ii) Sn + 2Ag⁺ → Sn²⁺ + 2Ag
 all species correct [1]
 accept equation with Sn⁴⁺
- (iii) Mn to Mn²⁺ need both species [1] electron loss **or** oxidation number increases [1]
- (iv) covered with oxide layer [1] makes it unreactive or protects or aluminium oxide unreactive [1]
- (b) (i) potassium has one valency electron [1]
- or loses one electron
 calcium has two valency electrons
 or loses two electrons
 [1]
 - (ii) potassium hydroxide → no reaction
 calcium hydroxide → calcium oxide and water
 ACCEPT metal oxide

| | Page 6 | | j | Mark Scheme | Syllabus | Paper |
|---|--------|------------|----------------------------|--|--------------------|---------------------|
| | | | | IGCSE – October/November 2008 | 0620 | 31 |
| | | [2] [2] | | | | |
| | | | [1] fc | or formulae of any TWO products | | |
| | | | | | | [Total: 17] |
| 7 | (a) | (i) | 35 cr 40 cr | | | [1] [1] |
| | | (ii) | form | s carbon monoxide | | [1] |
| | | | or ef | onous or toxic or lethal or prevents blood carrying of ffect on haemoglobin just harmful | oxygen | [1] |
| | (b) | (i) | | robutane or butyl chloride ber not required but if given must be 1, it must be ir | n correct position | [1] |
| | | (ii) | light | or UVor 200°C or lead tetraethyl | | [1] |
| | | (iii) | • | correct equation for example 2-chlorobutane ichlorobutane | | [1] |
| | (c) | (i) | CON | ect repeat unit ID continuation I(CH ₃)-CH ₂)- | | [1] [1] |
| | | (ii) | if nu | n-1-ol or butan-2-ol or butanol mber given then formula must correspond for secor ect position | nd mark and numbe | [1] r must be in |
| | | | CH ₃ - | ctural formula of above -CH ₂ -CH ₂ -CH ₂ OH or CH ₃ -CH(OH)-CH ₂ -CH ₃ -C ₄ H ₉ OH st mark not awarded then either formula will gain ma | ark [1] | [1] |
| | | (iii) | CH ₃ -NOT respe | EPT either formula for "butanol" $-CH(Cl)-CH_3 \text{ or } CH_3-CH_2-CH_2-Cl$ $-C_3H_7Cl$ onse must not include HCl uation given look at RHS only | | [1] |

[Total: 12]

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the October/November 2008 question paper

0620 CHEMISTRY

0620/32

Paper 32 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2008 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Second variant Mark Scheme

| Page 2 | Mark Scheme | Syllabus | Paper |
|--------|-------------------------------|----------|-------|
| | IGCSE – October/November 2008 | 0620 | 32 |

| 1 | ammonia chlorine "pop" with a lighted splint or burn with a pop or goes pop and extinguishes flame NOT glowing splint relights a glowing splint turns limewater milky/cloudy/chalky/white ACCEPT correct formulae | | | | | |
|---|---|--|-----------------------------|--|--|--|
| 2 | if ig if | Na : 1S correct ratio orrect charges e around S no symbols then must have correct key covalent only mark 1 nore electrons around sodium the response includes both a correct and an incorrect answer or not select correct one, mark = [0] | [1] [1] [1] | | | |
| | (b) (i) | positive ions or cations NOT atoms or cores or nuclei layers or lattice or regular pattern delocalised or free or mobile electrons or sea OR positive ions or cations NOT atoms or cores or nuclei attraction between ions and electrons delocalised or free or mobile electrons or sea the attraction/electrostatic bonding must be between ions and delocalised electrons, between cations and anions does not score ACCEPT bond if qualified e.g. electrostatic bond, etc. if moles or molecular cannot score cation mark delocalised/free/mobile electrons or electrons can move layers or ions or atoms or particles NB more flexible than 2(b)(i) can slip or move past each other or bonding non-directional | [1] [1] [1] [1] [1] [1] [1] | | | |

| | | IGCSE – October/November 2008 | 0620 | 32 |
|-------|------|--|--------------------|-------------------|
| (c) | (i) | tetrahedral 1Si : 4O bonded/surrounded, etc. 1O : 2 Si | | [1] [1] [1] |
| | | NOT molecules of oxygen, etc. NOT intermolecular forces ONLY tetrahedral can score for either of the above | | |
| | | Despite what the question states, ACCEPT a clear accuabove three points. | urate diagram whic | h shows the |
| | (ii) | hard high melting point or boiling point colourless (NOT clear) or shiny or translucent non/poor conductor (of electricity) brittle | | |
| | | insoluble any TWO NOT crystalline or strong | | [2] |
| | | | | [Total: 14] |
| 3 (a) | (i) | water or moisture ACCEPT salty water air or oxygen | | [1] [1] |
| | (ii) | galvanising or coat with zinc tin plate chromium plate nickel plate cobalt plate copper plate cover with aluminium anodic protection or sacrificial protection cathodic protection cover with plastic alloying (ignore any named metal) any TWO NOT just plate or electroplate need electroplate with su NOT oil | itable metal | [2] |
| (b) | (i) | ACCEPT both galvanising and sacrificial protection hydrogen or carbon or carbon monoxide or methane or more reactive metal NOT Group I | | [1] |

Mark Scheme

Syllabus

Paper

[2]

(ii) any correct equation

only error not balanced [1]

| Page 4 | | | Mark Scheme | Syllabus | Paper | | | |
|--------|-----|-------|-------------------------------|---|-------|-------------|--|--|
| | | | | IGCSE – October/November 2008 | 0620 | 32 | | |
| | (c) | (i) | 196 | | | [1] | | |
| | | (ii) | = 18 mark ONL | 36/196 × 100 = 18(.4)% ACCEPT 18 to nearest whole number mark e.c.f. to (c)(i) provided percentage not greater than 100% ONLY ACCEPT 36/answer (c)(i) × 100 otherwise [0] | | | | |
| | (d) | (i) | form | s carbon dioxide/carbon monoxide (which escapes |) | [1] | | |
| | | (ii) | | s silicon(IV) oxide or silicon oxide or silica CaO reacts with SiO ₂ | | [1] | | |
| | | | to for | rm slag or calcium silicate re an incorrect formula if a correct name given Si + O ₂ + CaO form slag | | [1] | | |
| | | | | | | [Total: 13] | | |
| 4 | (a) | (i) | | $_{5}$ COOH or C_{6} H $_{5}$ CO $_{2}$ H $_{5}$ C $_{7}$ H $_{6}$ O $_{2}$ / C_{6} H $_{6}$ COO | | [1] | | |
| | | (ii) | corre | um hydroxide + benzoic acid = sodium benzoate + ect spelling needed NOT benzenoate EPT correct symbol equation | water | [1] | | |
| | | (iii) | | um carbonate or oxide or hydrogencarbonate TWO · Na | | [2] | | |
| | (b) | (i) | 7.7% | 6 | | [1] | | |
| | | (ii) | | ny number: equal number ratio xample 1:1 or 6:6 | | [2] | | |
| | | (iii) | mole | irical formula is CH ecular formula is C ₆ H ₆ .c.f., award of marks not dependent on (ii) | | [1] [1] | | |
| | (c) | (i) | C ₆ H ₈ | $_{3}O_{6}$ | | [1] | | |
| | | (ii) | alcol NOT | on – carbon double bond or alkene hol or hydroxyl or hydroxy hydroxide oxide and alcohol = 0 | | [1] [1] | | |
| | | | | | | [Total: 12] | | |

| Page 5 | Mark Scheme | Syllabus | Paper |
|--------|-------------------------------|----------|-------|
| | IGCSE – October/November 2008 | 0620 | 32 |

- 5 (a) (i) $2H^+ + 2e \rightarrow H_2$ [1]
 - (ii) $2Cl^- 2e \rightarrow Cl_b$ or $2Cl^- \rightarrow Cl_b + 2e$ [1]
 - (iii) Na⁺ and OH⁻ are left [1] OR C*l*⁻ removed OH⁻ left

NB ions by name or formula essential NOT any reaction of Na or Na⁺

NOT Na⁺ and OH⁻ combine

(b) (i) sterilise/disinfect water or kill microbes/germs bacteria, etc.
 NOT just to make it safe to drink or purify it or clean it treat above as neutral they do not negate a correct response

(ii) ammonia **or** methanol **or** hydrogen chloride **or** margarine [1] **NOT** nylon

(iii) ester or triester or lipid [1] hydrolysis or saponification [1]

[Total: 7]

[1]

6 (a) (i)

| aqueous | tin | manganese | silver | zinc |
|-----------------------|-----|-----------|--------|------|
| solution | Sn | Mn | Ag | Zn |
| tin(II) nitrate | | R | NR | R |
| manganese(II) nitrate | NR | | NR | NR |
| silver(I) nitrate | R | R | | R |
| zinc nitrate | NR | R | NR | |

[3] for each row

ignore anything written in blank space

- (ii) Zn + 2AgNO₃ → Zn(NO₃)₂ + 2Ag
 all species correct [1]
 accept correct ionic equation
 Zn + 2Ag⁺ → Zn²⁺ + 2Ag [2]
- (iii) Sn²⁺ must be made clear that the oxidant is Sn²⁺ not Sn [1] it gains electrons **or** oxidation number decreases **or** it is reduced reason must relate to an oxidant **NB** not dependent on identifying Sn²⁺
- (iv) covered with oxide layer [1] makes it unreactive or protects or aluminium oxide unreactive [1]

| i age o | | ICCSE October/Nevember 2009 0620 | | 22 |
|-----------|-----------------------------------|--|---------------------|-------------|
| | | IGCSE – October/November 2008 | 0620 | 32 |
| (b) (i) | or lo | essium has one valency electron oses one electron ium has two valency electrons oses two electrons | | [1] |
| | or 10 | oses two electrons | | [1] |
| (ii) | calci | ssium hydroxide → no reaction ium hydroxide → calcium oxide and water CEPT metal oxide | | [1] [1] |
| (iii) | | $O_3 \rightarrow 2KNO_2 + O_2$ or formula of either product | | [2] |
| | | $(NO_3)_2 \rightarrow 2CaO + 4NO_2 + O_2$ or formulae of any TWO products | | [2] |
| | | | | [Total: 17] |
| ' (a) (i) | 20 cr 80 cr | | | [1] [1] |
| (ii) | • | s carbon monoxide | | [1] |
| | or et | onous or toxic or lethal or prevents blood carrying ffect on haemoglobin 「just harmful, etc. | g oxygen | [1] |
| (b) (i) | | robutane or butyl chloride ber not required but if given must be 1, it must be | in correct position | [1] |
| (ii) |) light | or UV or 200 °C or lead tetraethyl | | [1] |
| (iii) | | correct equation for example 2-chlorobutane | | |
| | | ichlorobutane t include HC <i>l</i> | | [1] |
| (c) (i) | | ect repeat unit | | [1] |
| | | ND continuation H(CH ₃)–CH ₂)– | | [1] |
| (ii) | if nu | oan-1-ol or propan-2-ol or propanol mber given then formula must correspond for sec | ond mark. | [1] |
| | struc | ber must be in correct position ctural formula of above –CH ₂ –CH ₂ –OH or CH ₃ –CH(OH)–CH ₃ | | [1] |
| | NOT if firs | 「C₃H₂OH st mark not awarded then either formula will gain n ept either formula for "propanol" in (i) On scoris both marks entered together not as ∣ | | , |
| (iii) | CH ₃ - NOT if eq | $-CH_2-CH_2-CH_2-Cl$ or $CH_3-CH_2-CH(Cl)-CH_3$ ΓC_4H_9Cl uation given look at RHS only | | [1] |

Mark Scheme

Syllabus

Paper

[Total: 12]

response must not include HCl