
Centre Number	Candidate Number	Name Man + Hit en
		Name Name SE INTERNATIONAL EXAMINATIONS ertificate of Secondary Education 0620/06
CHEMISTRY		0620/06
Paper 6 Alter	mative to Practical	October/November 2006
Candidates ansv No additional ma	wer on the Question Pap aterials required.	per. 1 hour
	e number and candidate	
'ou may use a pencil fo		

For Examiners Use		
1		
2		
3		
4		
5		
6		
Total		

This document consists of 13 printed pages and 3 blank pages.






(c)	Calculate the percentage of air used in the rusting of the iron.
	[2]
(d)	How would the results differ if pure oxygen was in the tube instead of air before rusting?
	[1]

For Examiner's Use


3 An investigation into the reaction of calcium with water was carried out using the apparatus below. The temperature of the water increased during the experiment.



The volume of hydrogen collected at one minute intervals was measured. Use the diagrams to record the volumes in the table.

time/minutes	syringe diagram	volume of gas/cm ³
0	10 20 30 40 50 60 70 80 90	
1		
2	10 20 30 40 50 60 70 80 90	
3	10 20 30 40 50 60 70 80 90	
4	10 20 30 40 50 60 70 80 90	
5	10 20 30 40 50 60 70 80 90	
6		

[2]



(a) Plot the results on the grid. Join all of the results with a smooth curve.

4 An investigation was carried out into the reactions of aqueous copper(II) sulphate with magnesium, iron and zinc.

Experiment 1

By using a measuring cylinder, 5 cm^3 of aqueous copper(II) sulphate was added to each of three test-tubes. The initial temperature of the solution was measured. Zinc powder was added to the first test-tube, iron powder to the second tube and magnesium powder to the third tube. The mixtures were stirred with the thermometer. All the observations were recorded and the maximum temperature reached measured.

(a) Use the thermometer diagrams to complete the results table.

Table of results	

metal added	temperature of solution/°C initial maximum		temperature difference/°C	observations
zinc		60 		moderate effervescence, solution paler, brown solid.
iron		40 40 35		little effervescence, brown solid.
magnesium	25 20	70 65		rapid effervescence, pops with lighted splint, brown solid.

[4]

(b) Use your results and observations to answer the following questions.

(i) Which metal is most reactive with aqueous copper(II) sulphate?

(c) The reactions of magnesium and zinc with aqueous copper(II) sulphate were investigated in more detail.

Experiment 2

By using a measuring cylinder 10 cm³ of aqueous copper(II) sulphate was poured into a polystyrene cup. The initial temperature of the solution was measured. A 1 g sample of magnesium powder was added to the cup and the temperature measured every 10 seconds for 1 minute.

Use the thermometer diagrams on **page 8** to complete the results table.

Experiment 3

Experiment 2 was repeated using zinc powder instead of magnesium.

Use the thermometer diagrams on **page 8** to complete the results table.


For
Examiner's
Use

Table of results

time/seconds		temperature/°C		
lime/seconds	Experiment 2		Experiment 3	
0			30 - 25 - 20	
10				
20	35 			
30	35 			
40	40 		75 70 - 65	
50	40		70 - 65 - 60	
60				

[6]

(d) Plot the results of both Experiments on the grid below. Draw two smooth line graphs. Clearly label the graphs.



(g)	Why is a polystyrene cup used instead of a glass container?	
		[1]
(h)	Suggest one improvement to the method in Experiment 2.	
		[1]

5 Two solids, F and G, were analysed. Solid F was an ammonium salt and solid G was a potassium salt.


The tests on **F** and **G** and some of the observations are in the following table.

Complete the observations in the table.

	tests	observations
and sha The sol	was added to distilled water aken to dissolve. ution was divided into 4 equal s in test-tubes.	
(a) (i)	The pH of the first portion of the solution was tested using Universal Indicator solution.	colour orange
(ii)	Aqueous sodium hydroxide was added to the second portion and heated gently.	рН 5
	The gas given off was tested with damp litmus paper.	[2]
(iii)	To the third portion of solution, was added dilute nitric acid and then aqueous lead(II) nitrate.	white precipitate
(iv)	To the fourth portion of solution, was added dilute nitric acid followed by aqueous silver nitrate.	white precipitate
(b) (i)	Solid G was dissolved in distilled water. The solution was divided into two test-tubes.	
(ii)	(a)(iii) was repeated using the first portion of the solution.	bright yellow precipitate
(iii)	(a)(iv) was repeated using the second portion of the solution.	pale yellow precipitate

(c)	What conclusion can be drawn from test (a)(i)?	
		[2]
(d)	Name the gas given off in (a)(ii) .	
		[1]
(e)	Identify solid F .	
		[1]
(f)	Identify solid G .	
		[1]

6 You are provided with a pot of paint as shown below.



BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.