

## **MARK SCHEME for the October/November 2012 series**

### **9700 BIOLOGY**

**9700/43**

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE AS/A LEVEL – October/November 2012</b>	<b>9700</b>	<b>43</b>

Mark scheme abbreviations:

<b>;</b>	separates marking points
<b>/</b>	alternative answers for the same point
<b>R</b>	reject
<b>A</b>	accept (for answers correctly cued by the question, or by extra guidance)
<b>AW</b>	alternative wording (where responses vary more than usual)
<b><u>underline</u></b>	actual word given must be used by candidate (grammatical variants excepted)
<b>max</b>	indicates the maximum number of marks that can be given
<b>ora</b>	or reverse argument
<b>mp</b>	marking point (with relevant number)
<b>ecf</b>	error carried forward
<b>I</b>	ignore
<b>AVP</b>	Alternative valid point (examples given as guidance)

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9700	43

- 1 (a) A - mitochondrion ;  
 B - post-synaptic membrane ;  
 C - myelin sheath / Schwann cell ; [3]
- (b) 1 produces ATP ; (1)  
 R produces energy  
*any two from*  
 2 (for) ACh production ;  
 3 (for) vesicle formation ;  
 4 (for) vesicle movement ;  
 5 (for) exocytosis / described ;  
 6 (for) functioning of ion pumps ;  
 R calcium ions (2 max) [3 max]
- (c) 1 fits into (membrane) receptors ;  
 2 not broken down (by enzymes) ;  
 3 (so) action potentials generated for a long time (in post-synaptic neurone) ;  
*ignore* ref to increased frequency of action potentials  
 4 AVP ; e.g. causes release of other transmitters / stimulant and depressant / variable response [2 max]
- [Total: 8]**
- 2 (a) 1 ref. sticky ends ;  
 2 GATC and CTAG ;  
 3 complementary bases (pairing) ;  
 4 A to T **and** C to G ;  
 5 H-bonds (to sticky ends of plasmid) ;  
 6 (gaps in) sugar-phosphate backbones sealed by (DNA) ligase ;  
 7 AVP ; e.g. formation of phosphodiester bonds / ref. terminal transferase [4 max]
- (b) (i) 1 *idea of* identifying bacteria that, are transformed / have taken up plasmid / have taken up ampicillin resistance gene ;  
 2 these bacteria have survived ;  
 3 these bacteria may contain pBR322 or recombinant plasmid / plasmids taken up may not contain human insulin gene ;  
 4 other bacteria have been killed ; [3 max]
- (ii) 1 (*Bam*HI) breaks the tetracycline resistance gene ;  
 2 (inserting human insulin gene) makes tetracycline resistance gene inactive ;  
 3 colonies that are ampicillin-resistant but **not** tetracycline-resistant have taken up recombinant plasmid / insulin gene ;  
 4 colonies that survive on, tetracycline / both ampicillin and tetracycline / plate **T**, have not taken up the recombinant plasmid / insulin gene ; [3 max]
- (iii) *Answer on Fig. 2.2*  
 left hand colony on plate **A** ; [1]

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9700	43

- (c) (i) 1 plasmids (easily) transferred between bacteria ;  
2 (bacteria of), same species / different species ;  
3 bacteria can acquire antibiotic resistance / renders antibiotic useless / AW ; [2 max]

(ii) *mark for gene and mark for how product detected*

- 1 gene for  $\beta$  galactosidase ;  
2 blue colour from X-gal medium ;  
**or**  
3 gene for  $\beta$  glucuronidase (GUS) ;  
4 produces product that is easily stained blue ;  
**or**  
5 gene for, GFP / other fluorescent product ;  
**R** fluorescent / fluorescence, gene  
6 fluorescence detected when present ;  
**or**  
7 other gene ;  
8 how detected ; [2 max]

**[Total: 15]**

- 3 (a) 1 globular ;  
2 ref. tertiary structure / 3D shape ;  
3 active site (because enzyme) ;  
4 outer amino acids with hydrophobic R groups (because in membrane) / AW ; [2 max]


- (b) 1 (penicillin) binds, rarely / briefly, with PBP2a ;  
*ignore* doesn't bind well  
2 (so) most PBP2a molecules not blocked ;  
3 (so) cell wall / cross links, can still be made (in presence of penicillin) ;  
4 penicillin is competitive inhibitor (of PBP) ;  
5 (so) reduces PBP enzyme activity ; [3 max]

- (c) 1 viruses have no (peptidoglycan) wall ;  
2 viruses have no, transpeptidase / glycoprotein peptidase ;  
3 viruses, have no cell structure / are not cells ;  
4 viruses have no metabolism ; [2 max]

**[Total: 7]**

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9700	43

4 (a)



[3]

- (b)
- 1 protein higher in whole grain flour **because** protein is in aleurone layer ;
  - 2 parts containing protein / aleurone layer, not removed (as in white flour) ;
  - 3 dietary fibre higher in whole grain flour **because** (most) fibre is in, pericarp / testa ;
  - 4 pericarp / testa, has not been removed (as in white flour) ;
  - 5 carbohydrate content lower in whole grain flour **because** outer parts not removed ;  
accept *ora* throughout [3 max]
- (c) (i) starch must be digested (to glucose) before it is absorbed / digestion of starch takes time ; [1]
- (ii)
- 1 amylose has 1–4 bonds / amylopectin has 1–4 bonds plus 1–6 bonds ;
  - 2 amylose, digested / broken down to glucose / acted on by amylase, more slowly ;
  - 3 because fewer sites for enzyme to work on / AW ;  
accept *ora* for mp2 and mp3 [2 max]

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9700	43

- (d) (i) 1 increasing intake (of whole cereal grains) decreases risk (of developing type II diabetes);  
2 use of figures supporting this relationship ;  
3 not all values fit the trend / reference to this not being a linear effect ;  
4 reference to higher risk at 19.0 – 24.5 intake ; [3 max]
- (ii) 1 idea that the risk of 1.00 for each food group is not the same risk ;  
2 no info on size of servings / no indications that same units used for each group ;  
3 intervals of range of intake not consistent – different intervals may give different results ; [2 max]
- (iii) 1 fruits contain, sugars / glucose / fructose ;  
2 sugar has a high GI ; [2]

[Total: 16]

- 5 (a) 1 ref. to suitable container e.g. dish  
**or**  
ref. suitable medium ;  
2 ref. to addition of, sperm / semen, to oocytes ; [2]  
**A ICSI**

- (b) *advantage*  
better chance of survival / more certain of getting a good-quality embryo / better chance of implantation ;  
*disadvantage*  
may be difficult to keep embryos alive for this time / embryos may become less viable / less chance of implantation ; [2]  
*only allow one mark for ref. to implantation*

- (c) (i) 1 higher % of pregnancies than the other methods ;  
2 35.1 % versus 22 .1 % **or** 35.1 % versus 34.6 % ;  
3 little difference in the success rate of single top quality embryo transfer compared to multiple embryo transfer ;  
4 multiple embryos increases risk of problems during pregnancy / birth ; [3 max]
- (ii) 1 could lead to selection of features desired by parents / society  
**or** less chance of a child being born with features seen as undesirable ;  
2 ref. to discarding other embryos ; [1 max]

[Total: 8]

Page 7	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9700	43

- 6 (a) (i) *accept answers in a genetic diagram where genotypes are linked to phenotypes*
- 1 agouti allele /  $C^a$ , dominant to black allele /  $C^b$  ; **ora**
  - 2 black parents homozygous recessive ;
  - 3 agouti parents heterozygous **or** homozygous ; [2 max]
- (ii) *accept answers in a genetic diagram where genotypes are linked to phenotypes*
- 1 yellow allele /  $C^y$ , dominant to, black allele /  $C^b$  ;
  - 2 ref. to modified 3:1 ;
  - 3 (homozygous) genotype  $C^y C^y$ , lethal / does not survive ; [2 max]
- (iii) *accept answers in a genetic diagram where genotypes are linked to phenotypes*
- 1 yellow allele /  $C^y$ , dominant to **all** others ;
  - 2 agouti /  $C^a$  **or** black and tan /  $C^{bt}$ , allele, dominant to black allele ;  
**A** black allele recessive to all other alleles
  - 3 yellow mice all heterozygous (must be stated) ; [2 max]
- (b) 1 cross (black and tan mouse) with, black mouse / homozygous recessive mouse /  $C^b C^b$  ;
- 2 if **all** offspring black and tan then parent,  $C^{bt} C^{bt}$  / homozygous ;
  - 3 if some offspring are black (and some are black and tan) then parent,  
 $C^{bt} C^*$  / heterozygous ; [2 max]

[Total: 8]

- 7 (a) 1 *idea of genetic variation* ;
- 2 increased heterozygosity / decreased homozygosity ;
  - 3 hybrid vigour / decreased inbreeding depression ;
  - 4 able to adapt to changing conditions ;
  - 5 *idea of some individuals surviving* ;
  - 6 AVP ; e.g. reduced risk of expression of harmful recessive alleles [3 max]

- (b) (i) *most affected*  
almond, because, 100% / all / only, pollinated by honey bee ;  
*least affected*  
orange, because only 25% pollinated by honey bee / 75% pollinated by other methods [2]

- (ii) *any three from*
- 1 parasites / mites / viruses / bacteria ;  
**A** disease
  - 2 detail of climate change ; e.g. temperature change
  - 3 pollution qualified ; e.g. increased use of pesticides / increased sulfur dioxide concentration in air
  - 4 inbreeding ;
  - 5 competition for food / food shortage ;
  - 6 increase in predator numbers ;
  - 7 AVP ; e.g. ref. killer bees / plant monoculture provides limited nutrition [3 max]

[Total: 8]

Page 8	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9700	43

- 8 (a) (i) cytoplasm / cytosol ; [1]
- (ii) 1 NAD regenerated ;  
2 so glycolysis can continue ;  
3 to produce ATP ; [2 max]
- (iii) lactate dehydrogenase ; [1]
- (iv) *reaction* - condensation / polymerisation ;  
*bond* - glycosidic ; [2]
- (b) *in yeast*
- 1 decarboxylation / CO<sub>2</sub> removed ;  
2 ethanal (as intermediate step) ;  
3 ethanol produced ;  
4 two steps (from pyruvate) ;  
5 ethanol dehydrogenase ;  
6 not a reversible reaction / ethanol cannot be converted back to pyruvate ;  
7 *idea of process less energy efficient ;*  
*allow **ora** for mp1, mp4, mp5, mp6 and mp7* [4 max]
- (c) (i) carbon dioxide produced divided by oxygen consumed ;  
volume / number of moles (of both gases) ; [2]
- (ii) carbohydrate = 1.0 ;  
lipid = 0.7 ; [2]
- (iii) increase / go above one / infinity ; [1]
- [Total: 15]**
- 9 (a) 1 occur during meiosis I ;  
*crossing over*  
2 between non-sister chromatids ;  
3 of, (a pair of) homologous chromosomes / a bivalent ;  
4 in prophase 1 ;  
5 at chiasma(ta) ;  
6 exchange of genetic material / AW ;  
*R genes unqualified*  
7 linkage groups broken / AW ;  
8 new combination of alleles (within each chromosome) ;  
*independent assortment*  
9 of homologous chromosomes pairs / bivalents ;  
10 each pair lines up independently of others ;  
11 line up on equator ;  
12 (during) metaphase 1 ;  
13 results in gametes that are genetically unique / AW ; [9 max]



(b)

	<i>artificial selection</i>		<i>natural selection</i>
14	selection (pressure by) humans	or	environmental selection pressure ;
15	genetic diversity lowered	or	genetic diversity remains high ;
16	inbreeding common	or	outbreeding common ;
17	loss of vigour / inbreeding depression	or	increased vigour / less chance of inbreeding depression ;
18	increased homozygosity / decreased heterozygosity	or	decreased homozygosity / increased heterozygosity ;
19	no isolation mechanisms operating	or	isolation mechanisms do operate ;
20	(usually) faster	or	(usually) slower ;
21	selected feature for human benefit	or	selected feature for organism's benefit ;
22	not for, survival / evolution	or	promotes, survival / evolution ;

[6 max]

[Total: 15]

- 10 (a)
- 1 PII absorbs light ;
  - 2 enzyme (in PII) involved ;
  - 3 to break down water / AW ;
  - 4  $2\text{H}_2\text{O} \longrightarrow 4\text{H}^+ + 4\text{e}^- + \text{O}_2$  ;
  - 5 oxygen is produced ;
  - 6 used by cells for (aerobic) respiration ;
  - 7 or released (out of plant) through stomata ;
  - 8 protons used to reduce NADP ;
  - 9 with electrons from PI ;
  - 10 reduced NADP used in, light independent stage / Calvin cycle ;
  - 11 to convert GP to TP ;
  - 12 electrons also used in ETC ;
  - 13 to release energy for photophosphorylation ;
  - 14 to produce ATP ;
  - 15 electrons (from PII) go to PI ;
  - 16 ref. re-stabilise PI ;

[10 max]

<b>Page 10</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE AS/A LEVEL – October/November 2012</b>	<b>9700</b>	<b>43</b>

- (b) 16 gibberellin is a, plant growth regulator / plant hormone / plant growth substance ;  
17 stimulates cell division ;  
18 stimulates cell elongation ;  
19 detail of cell elongation ; e.g. changes plasticity of cell wall  
20 plant grows tall ;  
21 apply gibberellin to dwarf plants and they grow taller / gibberellin promotes bolting of some rosette plants ;  
22 ref. inactive and active forms ;  
23 dwarf plants, lack active form / have inactive form, of gibberellin ;  
24 (dominant) allele causes synthesis of enzyme ;  
25 (enzyme) catalyses the production of the active form of gibberellin ;  
26 recessive allele only inactive form of gibberellin formed / dominant allele results in active form of gibberellins ;  
27 AVP ; e.g. ref. to different forms of gibberellins / there is interaction between / gibberellin and other plant growth regulators [5 max]

**[Total: 15]**