CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2013 series

9700 BIOLOGY

9700/51

Paper 5 (Planning, Analysis and Evaluation), maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Mark schemes abbreviations:

; separates marking points

I alternatives answers for the same point

R do not allow

A allow (for answers correctly cued by the question, or guidance for examiners)

AW alternative wording (where responses vary more than usual)

<u>underline</u> actual word given must be used by candidate (grammatical variants excepted)

max indicates the maximum number of marks that can be given

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Extra guidance	Mark	АО
1 (a) (i)	independent: concentration of GA; dependent: ref. to the starch free, area / zone (around grains);	ignore amount / quantity A diameter / radius / size A clear zone / brown zone / digested starch ignore ref. to amylase activity	[2]	Р
(ii)	8 of: independent variable	A seed / fruit / maize for any mark point in (a) ignore any ref. to time between soaking and putting maize onto agar plate		
	 ref. to method of diluting the (3 mmol dm⁻³) GA to give a minimum of (any) 5 dilutions; 	A series / serial / proportional / simple, dilution as method or description of method		
	2. ref. to concentrations (other than 0) that fall in the range 3 mmol dm ⁻³ to any value above zero with units;	2. minimum is 2 values that are not higher than $3\text{mmol}\text{dm}^{-3}$ and are above zero $1\text{mmol}\text{dm}^{-3} = 1000\mu\text{mol}\text{dm}^{-3}$ $1\mu\text{mol}\text{dm}^{-3} = 0.001\text{mmol}\text{dm}^{-3}$		
	3. <i>ref. to</i> soaking grains in GA solutions for, min 24 hours / max 72 hours ;	 ignore if the range of GA is incorporated into agar plates If pre-soaked in water for 24h and then in GA, must be minimum of 1 hour GA 		
	4. ref. to a control using (distilled / deionised) water;	4. A description – 'to allow comparison' / AW		
	5. ref. to stated incubation temperature;	 5. e.g. any one temp in the range 15–35°C. A room temperature. R body temperature / 37°C 		

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Extra guidance	Mark	АО
	dependent variable			
	6. ref. to a suitable method of measuring the starch free area around the grains; standardising variables (max 4):	 6. e.g. trace outline onto a grid / transparent grid / photograph and put on grid / cut out clear zone and put on grid, and count squares. OR use suitable apparatus to measure, diameter / radius, directly or from a grid and calculate area using πr² ignore metre ruler. A e.g. ruler / callipers / micrometer / eyepiece graticule 		
	7. ref. to using, same / stated number of, grains / halves, for each concentration;	7. A 'one / a' as same number ignore 'same size of grains / same amount of grains. A quantity for number		
	8. ref. to, stated / same, volume of each soaking solution / GA;	8. A idea of: all, submerged / covered		
	9. ref. to method of keeping same incubation temperature;	9. e.g. water bath / incubator / temperature controlled room / thermostatically controlled environment / environmental chamber / propagator / thermostat ignore air conditioning		
	10. ref. to one stated time for incubation (of enzyme);	10. A any value in the range 24–72 hours. Actual value must be statedA left on, plate / agar as AW for incubate		
	11. ref. to, same / stated, concentration of starch in agar;	11. A 'known' time		
	12. ref. to same, volume / depth, of agar (in the Petri dishes);			
	13. covering to prevent, evaporation / contamination;			

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Extra guidance	Mark	AO
	safety: 14. ref. to, low risk investigation / any suitable safety precaution;	14. e.g. cutting away from hands / using tile for cutting. ignore gloves for cutting. ref. to allergy / irritation, and, gloves / mask / eye protection R no risk		
	reliability: 15. ref. to replicates, and mean / to identify or eliminate anomalies;	 15. must be a minimum of 3 (data sets), allow as, original and 2 more / several A outliers for anomalies A average for mean 	[8]	М
(b)	x-axis: concentration of GA and mmol dm ⁻³ or μmol dm ⁻³ ;	A mmol or μmol, per dm³ or mmol / dm³ or μmol / dm³		
	y-axis: (mean) area of, clear zone / starch digested and mm ² ;	A (mean) diameter of clear zone / mm A cm² / cm A activity of amylase and arbitrary units / au labels correct – no / incorrect, unit(s) = 1 max axes reversed, correct for labels and units = 1 max	[2]	D
(c) (i)	3 of: 1. idea of areas irregular in shape;	ignore ref. to replicates / repeats 1. A in terms of different diameters A in general terms like 'difficult to tell actual size'		
	2. idea of edges of areas difficult to see;	A in terms of edges overlapping or edges blurred / brown merging into blue / AW		
	3. not sure how much GA embryo is also producing;	3. A grain / fruit / seed for embryo		
	4. may be other, enzymes / chemicals, that hydrolyse the starch;			

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Ext	tra guidance	Mark	АО
	5. idea of: no way of quantifying amylase (leaving grain);	5.	A in terms of, not diffusing out / used inside / ref. to different surface area of (cut) grain / size of grain only in context of amylase not leaving grain or having different amounts of amylase		
	6. amylase / enzyme, may come from contamination;	6.	A coming from fungi / bacteria / coat of grain		
	7. idea of: no method of quantifying the starch disappearance;	7.	idea of: do not know of the number of molecules of starch broken down. Look in context of the test being qualitative rather than quantitative. NOT rate ideas		
	8. pH may vary / AW ;		14040	[3]	Е
(ii)	$\begin{tabular}{ll} \it method: any 1 \times 2 \ of: \\ \it 1. & stated \ better \ method \ of \ measuring, \ (area \ / \ diameter) \ ; \\ \it some \ brief \ detail \ of \ stated \ method \ ; \\ \end{tabular}$	1.	e.g. use graticule on low power of microscope = 2 marks		
	 if overlap: separate plates / fewer grains per plate / larger plates; able to distinguish edges more clearly / AW; if 'blur': really cannot be improved = 2 marks;; or do large sample; minimises, effect / error / AW; 				
	3. idea of: separating the embryo from the grains (after soaking); idea of: measure area, without embryo present / with embryo on its own;				
	4. <i>idea of:</i> extracting the amylase (from grain); <i>idea of:</i> repeat with just the extract and compare;				

Page 7	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Extra guidance	Mark	АО
	5. if in term of how much out / used up: idea of: extracting the amylase (from grain); idea of: repeat with just the extract and compare; if in terms of different surface areas: idea of: measuring (surface) area; idea of: either finding similar area or calculating per unit area;	5. A remove endosperm repeat using embryo / AW		
	6. <i>idea of:</i> sterilising / disinfecting (grain / agar / dish /AW); detail of possible method;	 detail e.g. (named) disinfectant or sterile technique to, make plates / make up agar / handle grains or keep plate covered. 		
	7. <i>idea of:</i> putting known concentrations of amylase on starch agar; producing a calibration curve from areas measured; 8. use a buffer;			
	to make up the agar (plates);		[2]	М
		Total:	[17]	

Page 8	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Extra guidance	Mark	АО
2 (a)	 3 of: 1. anaerobic bacteria increase with depth and aerobic bacteria decrease with depth; 2. below 1.5–2.0 m aerobic stay approximately the same and anaerobic continues to increase(slowly); 	A general point – but must relate to idea of depth		
	3. <i>idea of:</i> oxygen content of soil decreases with depth;4. <i>idea that:</i> anaerobic bacteria replace the aerobic bacteria as	3. A oxygen content decreases with time4. A replacement in terms of succession /		
	oxygen reduces / (most) aerobic unable to survive in low oxygen;	competition	[3]	С
(b) (i)	less activity of, dehydrogenase / enzyme ;	A less <u>aerobic</u> respiration R less dehydrogenase	[1]	E
(ii)	 2 of: temperature; pH; idea of: time that determination was run for; (soil) moisture / water; substrate (in soil); 	ignore ref. to oxygen5. A (named) nutrients (in soil)	[2]	P
(iii)		A heated / boiled / disinfected / sterilised / irradiated / AW, soil A (soil) with, dead / killed bacteria / denatured enzyme / enzyme removed A ref. to add an enzyme inhibitor A remove the bacteria from soil R killed enzyme / replace with (named) inert material	[1]	P

Page 9	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9700	51

Question	Expected answer	Extra guidance	Mark	AO
(c) (i)	ref. to wide / long / big / AW, error bars, that indicate results are, less / not, reliable; ref. to data;	A reverse argument A suitable AW for wide e.g. 6 years least reliable or 6 months / 3 years most reliable or more reliable in B than A at 6 years	[2]	D
(ii)	6 months and 3 year , 6 months and 6 years both underlined / ringed; error bars do not overlap;	A error bars do not cross / AW	[2]	D
(d)	idea of: graph of, dehydrogenase / enzyme, activity, against number of bacteria per gram (plotted); idea of: dehydrogenase / enzyme, activity located on graph and number of bacteria read from scale;	 ignore orientation of graph. A if shown on a graph for 2 marks A idea of: comparing data against table to look for 'closest approximation' = 1 mark max ignore compare unqualified 	[2]	D
		Total:	[13]	