

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME				
		CANDIDATE NUMBER		
CHEMISTRY		9701/22		
Paper 2 Structured Questions AS Core		October/November 2009		
Candidates and	swer on the Question Paper.	1 hour 15 minutes		

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE ON ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units. A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part question. At the end of the examination, fasten all your work securely together.

For Examiner's Use		
1		
2		
3		
4		
5		
Total		

This document consists of **11** printed pages and **1** blank page.



Answer **all** the questions in the spaces provided.

1 The elements carbon and silicon are both in Group IV of the Periodic Table. Carbon is the second most abundant element by mass in the human body and silicon is the second most common element in the Earth's crust.

Carbon and silicon each form an oxide of general formula XO_2 . At room temperature, CO_2 is a gas while SiO₂ is a solid with a high melting point.

(a) Briefly explain, in terms of the chemical bonds and intermolecular forces present in **each** compound, why CO₂ is a gas and SiO₂ is a solid at room temperature.

.....[3]

(b) Draw a simple diagram to show the structure of SiO₂. Your diagram should contain at least **two** silicon atoms **and** show clearly how many bonds each atom forms.

[2]

For Examiner's Use CO_2 does not behave as an ideal gas. For Examiner's Use (c) (i) State the basic assumptions of the kinetic theory as applied to an ideal gas. (ii) Suggest **one** reason why CO_2 does not behave as an ideal gas. [5] Carbon exists in a number of forms, one of which is a conductor of electricity and one of which is a non-conductor of electricity. Silicon is the main component of most semi-conductors. (d) Graphite is the form of carbon that is a conductor of electricity. Give a simple explanation for this property. When carbon and silicon(IV) oxide are heated together at about 2000 °C, silicon carbide, SiC, is formed. Silicon carbide is a hard material which is widely used as an abrasive and in ceramics. (e) (i) Construct an equation for the reaction of carbon and silicon(IV) oxide. SiC has a similar structure to one of the common forms of carbon. (ii) Which form is this? Give a reason for your answer. form reason [2] [Total: 13]

3

2 The elements of the third period of the Periodic Table form chlorides of general formula ECl_x where *E* represents the element. These chlorides show a variation in oxidation number from sodium to sulfur.

For Examiner's Use

(a) (i) Use the information given to complete the table below.

formula of chloride	NaC1	MgCl ₂	AlCl ₃	SiCl ₄	PCl ₃	SCl ₂
oxidation number of element in the chloride						

(ii) By considering the electron configurations of the elements, explain the variation in oxidation number in the chlorides from Na to A*l* and from Si to S.

Na to Al	
Si to S	
	[5]

Sodium hydride, NaH, is a colourless crystalline solid which melts at 800 °C and has the same crystal structure as sodium chloride which has a melting point of 808 °C. When molten sodium chloride is electrolysed using graphite electrodes, a shiny deposit, **D**, forms on the cathode and a greenish-yellow gas is evolved from the anode. When molten sodium hydride is electrolysed, under suitable conditions using graphite electrodes, the same shiny deposit **D** is formed on the cathode and a colourless gas, **G**, is evolved from the anode.

(b) (i) Describe with the aid of a diagram the bonding in a sodium chloride crystal.

(ii) Suggest the type of bonding that is present in sodium hydride.

.....

(iii) What is the oxidation number of hydrogen in sodium hydride?

.....

For Examiner's Use

(v) The metals magnesium and aluminium form hydrides with formulae MgH₂ and AlH_3 . The non-metals phosphorus and sulfur form hydrides with formulae PH₃ and H₂S.

By considering their positions in the Periodic Table, suggest oxidation numbers for these four elements in their hydrides.

compound	MgH_2	AlH3	PH_3	H ₂ S
oxidation number of element in the hydride				

[8]

At room temperature, the chlorides of sodium, magnesium and aluminium are all solids which dissolve in water.

The hydrides of sodium, magnesium and aluminium are also solids which react with water with the rapid evolution of the **same** colourless gas **G** in each case.

(c) (i) What is the pH of the solutions formed when separate samples of sodium chloride, magnesium chloride, and aluminium chloride are dissolved in water?

chloride	sodium	magnesium	aluminium
рН			

(ii) Suggest an equation for the reaction between sodium hydride and water.

.....

(iii) Suggest a value for the pH of the solution formed in (ii).

.....

[4]

At room temperature, the chlorides of silicon, phosphorus and sulfur are all low melting point solids or low boiling point liquids that can be seen to react with water.

(d) (i) Suggest what type of bonding is present in sulfur dichloride, SCl_2 .

.....

(ii) Write a balanced equation for the reaction between the chloride of silicon, $SiCl_4$, and water.

.....[2]

[Total: 19]

3 One method of making 1-bromobutane in the laboratory is described below.

		Use
Stage 1	Place 35 g of powdered sodium bromide, 30 cm ³ of water, and 25 cm ³ (20 g) of butan-1-ol, in a 250 cm ³ two necked flask fitted with a tap funnel and reflux condenser.	
Stage 2	Concentrated sulfuric acid (25 cm ³) is then placed in the tap funnel and added drop by drop to the reagents in the flask, keeping the contents well shaken and cooled occasionally in an ice-water bath.	

(a) The overall reaction may be considered to take place in two stages. In the first stage the inorganic reagents react together to form HBr. In the second stage, the organic reagent reacts with the HBr that is formed in the first stage.

Write an equation for **each** of these stages.

(b) In this preparation, by using the amounts given above, **one** of the reagents, sodium bromide or butan-1-ol, will be present in an excess.

Use your equations in (a) and the data above to determine, by calculation, which reagent is in an excess.

[2]

For Examiner's

(c) In a laboratory preparation of 1-bromobutane, when 15.4g of butan-1-ol was used, 22.5g of 1-bromobutane was obtained after purification.

Calculate the yield of 1-bromobutane as a percentage of the theoretical maximum yield.

[2]

(d) When the concentrated sulfuric acid is added to the reaction mixture (stage 2), unless the temperature is controlled carefully, the acid may react with either of the original reactants (sodium bromide or butan-1-ol) to give at least two by-products, one of which is inorganic and the other organic.

For Examiner's Use

What inorganic and organic by-products may be formed?

In **each** case, identify **one** by-product and state the role of the concentrated sulfuric acid in the formation of this by-product.

inorganic by-product

role of conc. H_2SO_4

organic by-product

role of conc. H₂SO₄[4]

[Total: 10]

 4 (a) Complete the following reaction scheme which starts with 1-bromobutane. In each empty box, write the structural formula of the organic compound that would be formed.





[2]

[Total: 6]

- **5** The fermentation of starch or molasses using the bacterium *Clostridium acetobutylicum*, produces a mixture of propanone and butan-1-ol.
 - (a) Give the reagent(s) and state what would be observed when **one** test is carried out to confirm the presence of propanone in a mixture of propanone and butan-1-ol.

For Examiner's Use

- (c) Draw the following formulae.
 - (i) the structural formula of pentan-1-ol

(ii) the displayed formula of pentan-2-ol

(iii) the skeletal formula of pentan-3-ol



(e) (i) Draw the structural formula of the derivative of propanol that has the molecular formula $C_5H_{12}O$.

(ii) Draw the structural formula of the organic compound that will be present when the derivative of propanol you have given in (i) is heated under reflux with acidified potassium dichromate(VI).

[2]

[3]

[Total: 12]

11

For

Examiner's Use

(d) Identify this alcohol and give the structural formula of each alkene.

formulae are formed.

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.