



Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME											
CENTRE NUMBER								ANDIDATE IUMBER			
CHEMISTRY										970)1/32
Paper 3 Advance	ced Prac	ctical Ski	lls 2						May/	June	2016
										2 h	ours
Candidates ans	wer on t	he Ques	tion Pa	aper.							
Additional Mater	rials:	As liste	ed in th	ne Co	onfidentia	al Instructions	8				

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11. A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

Session	า
Laborato	ry

For Examiner's Use				
1				
2				
3				
Total				

This document consists of 12 printed pages.



1 In Questions 1 and 2 you will determine the relative atomic mass, A_r , of magnesium by two different methods.

In the first method you will collect and measure the volume of gas given off in the reaction between a known mass of magnesium and a known amount of dilute sulfuric acid. The acid will be in excess.

$$Mg(s) + H_2SO_4(aq) \rightarrow MgSO_4(aq) + H_2(g)$$


FB 1 is a strip of magnesium ribbon, Mg.

FB 2 is 1.00 mol dm⁻³ sulfuric acid, H₂SO₄.

(a) Method

Read through the whole method before starting any practical work.

The diagrams below may help you in setting up your apparatus.



- Fill the tub with water to a depth of about 5 cm.
- Fill the 250 cm³ measuring cylinder **completely** with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so the open end is in the water just above the base of the tub.
- Pipette 25.0 cm³ of FB 2 into the reaction flask labelled X.
- Check that the bung fits tightly in the neck of flask **X**, clamp flask **X**, and place the end of the delivery tube into the inverted 250 cm³ measuring cylinder.
- Weigh the magnesium ribbon, **FB 1**, and record the mass in the space below.
- Remove the bung from the neck of the flask. Add the magnesium ribbon, **FB 1**, into the acid in the flask and replace the bung **immediately**. Remove the flask from the clamp and swirl it to mix the contents. Swirl the flask occasionally until no more gas is evolved. Replace the flask in the clamp.
- When no more gas is collected, measure and record the final volume of gas in the measuring cylinder in the space below.

KEEP THE CONTENTS OF THE CONICAL FLASK X FOR USE IN QUESTION 2.

Results

1	(b)	Cal	cul	lati	ons
۱		, vai	ıvu	ıatı	OHIG

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of gas collected in the measuring cylinder. [Assume that 1 mole of gas occupies 24.0 dm³ under these conditions.]

moles of gas = mol

(ii) Use your answer to (i) and the mass of magnesium, **FB 1**, recorded in (a) to calculate the relative atomic mass, A_r , of magnesium.

 A_{r} of magnesium =[3]

[Total: 5]

2 You will determine the amount of sulfuric acid remaining in flask **X** after the reaction with magnesium in **Question 1**. You will do this by titration with sodium hydroxide of known concentration.

$$2NaOH(aq) + H2SO4(aq) \rightarrow Na2SO4(aq) + 2H2O(I)$$

FB 3 is 0.150 mol dm⁻³ sodium hydroxide, NaOH. bromophenol blue indicator

(a) Method

- Transfer all the contents of flask **X** from **Question 1** into the 250 cm³ volumetric flask.
- Rinse flask **X** with distilled water and add the washings to the volumetric flask. Add distilled water up to the mark. Stopper the volumetric flask and mix the contents thoroughly.
- Label this solution FB 4.
- Rinse the pipette and use it to transfer 25.0 cm³ of **FB 4** into the conical flask.
- Add about 10 drops of bromophenol blue to the conical flask.
- Fill the burette with FB 3.
- Perform a rough titration and record your burette readings in the space below. The end point is reached when the solution becomes a permanent blue-violet colour.

The rough titre	is		cm^3
-----------------	----	--	--------

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Record, in a suitable form below, all of your burette readings and the volume of FB 3
 added in each accurate titration.
- Make certain any recorded results show the precision of your practical work.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

(b) From your accurate titration results, obtain a suitable value for the volume of **FB 3** to be used in your calculations. Show clearly how you obtained this value.

25.0 cm³ of **FB 4** required cm³ of **FB 3**. [1]

(C)	Ca	lc	ul	af	ic	n	s
۱		•	Оũ	ľ	uı	a		/!!	J

Show y	our working	and appropria	ate significan	t figures in	the final a	answer to e	ach step	of your
calculat	tions							

(i) Calculate the number of moles of sodium hydroxide, NaOH, in the volume of **FB 3** you calculated in **(b)**.

moles of NaOH = mol

(ii) Use your answer to (i) to calculate the number of moles of sulfuric acid present in the 25.0 cm³ of **FB 4** pipetted in (a).

moles of $H_2SO_4 = \dots mol$

(iii) Use your answer to (ii) to calculate the number of moles of sulfuric acid, H₂SO₄, remaining in flask **X** after the reaction in **1(a)**.

moles of H_2SO_4 remaining from **1(a)** = mol

(iv) Use the relevant information on page 2 to calculate the number of moles of sulfuric acid, H₂SO₄, pipetted into reaction flask **X** in **1(a)**.

moles of H_2SO_4 pipetted into flask **X** = mol

(v)	Use your answers to (iii) and (iv) to calculate the number of moles of sulfuric acid which reacted with the magnesium in flask X .				
(vi)		er to (v) and	d the mass of ma		sk X = mol 1(a) to calculate the relative
	atomic mass, A	,, of magne:	sium.		
				$A_{\scriptscriptstyle{ m r}}$ of ma	gnesium =[6]
(d) (i)	A student, who A_r of magnesium			in Questions 1 a	nd 2 correctly, calculated the
			Question 1	Question 2	
		A _r Mg	20.8	22.5	
	procedure is les	ss accurate.		rce of inaccuracy	12 to deduce which practical and explain one change the
		is le	ess accurate		
	source of inacc	uracy			
	improvement				
(ii)	Use the A _r of m the student's va			odic Table to calc	ulate the percentage error in
					[3]

[Total: 17]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs. Marks are **not** given for chemical equations.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given. (a) (i) FB 5 is a solid element and FB 6 is a solid compound containing one cation and one anion. Carry out the following tests and record your observations.

test	observations
Place a small spatula measure of FB 5 in a boiling tube, add a 1 cm depth of dilute hydrochloric acid and warm the contents of the tube gently.	
Place small spatula measures of FB 5 and FB 6 in a single boiling tube. Use a test-tube holder to hold the tube. Add a 2 cm depth of aqueous sodium hydroxide. CARE	
Place a spatula measure of FB 6 in a test-tube. Add a 3 cm depth of distilled water to form a solution for the following two tests.	
To a 1 cm depth of aqueous FB 6 in a test-tube add a 1 cm depth of dilute hydrochloric acid.	
To a 1 cm depth of aqueous FB 6 in a test-tube add aqueous sodium hydroxide.	

Suggest the identities of FB 5 and FB 6 from your observations. Re observations in your answers.	fer to the relevant
FB 5:	
reason(s)	
FB 6: cation reason(s)	
anion reason(s)	

[9]

© UCLES 2016 9701/32/M/J/16

(ii)

- (b) (i) FB 7 and FB 8 are solutions which contain different anions. These may be carbonate, halide, sulfate or sulfite. You are to devise tests to identify the two anions present. Record in a suitable table below:
 - the reagent(s) you use for each test,
 - the observations you make on carrying out the test,
 - the conclusion you make from the result of the test.

(ii) The cation in either FB 7 or FB 8 is a transition metal ion. Carry out the following test to identify this cation and record your observations.

toot	observations				
test	FB 7	FB 8			
To a 1 cm depth of solution in a test-tube add aqueous sodium hydroxide.					

FB contains the transition metal ion

[9]

[Total: 18]

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

	reaction with									
ion	NaOH(aq)	NH ₃ (aq)								
aluminium, A <i>l</i> ³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess								
ammonium, NH₄⁺(aq)	no ppt. ammonia produced on heating									
barium, Ba ²⁺ (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.								
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.								
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess								
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution								
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess								
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess								
magnesium, Mg²+(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess								
manganese(II), Mn²+(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess								
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess								

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, C <i>l</i> ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and A <i>l</i> foil
nitrite, NO ₂ ⁻ (aq)	NH_3 liberated on heating with $OH^-(aq)$ and Al foil; NO liberated by dilute acids (colourless $NO \rightarrow$ (pale) brown NO_2 in air)
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO ₂)
chlorine, Cl ₂	bleaches damp litmus paper
hydrogen, H ₂	"pops" with a lighted splint
oxygen, O ₂	relights a glowing splint

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

The Periodic Table of Elements

Group																	
1	2											13	14	15	16	17	18
Key 1 H Hydrogen 1.0												2 He helium 4.0					
3	4			atomic numbe				_				5	6	7	8	9	10
Li	Be		atomic symbol									В	С	N	0	F	Ne
lithium 6.9	beryllium 9.0		rela	name ative atomic m	ass							boron 10.8	carbon 12.0	nitrogen 14.0	oxygen 16.0	fluorine 19.0	neon 20.2
11	12					J						13	14	15	16	17	18
Na	Mg											Αl	Si	P	S	C1	Ar
sodium 23.0	magnesium 24.3	3	4	5	6	7	8	9	10	11	12	aluminium 27.0	silicon 28.1	phosphorus 31.0	sulfur 32.1	chlorine 35.5	argon 39.9
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
potassium 39.1	calcium 40.1	scandium 45.0	titanium 47.9	vanadium 50.9	chromium 52.0	manganese 54.9	iron 55.8	cobalt 58.9	nickel 58.7	copper 63.5	zinc 65.4	gallium 69.7	germanium 72.6	arsenic 74.9	selenium 79.0	bromine 79.9	krypton 83.8
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	Υ	Zr	Nb	Мо	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Те	I	Xe
rubidium 85.5	strontium 87.6	yttrium 88.9	zirconium 91.2	niobium 92.9	molybdenum 95.9	technetium -	ruthenium 101.1	rhodium 102.9	palladium 106.4	silver 107.9	cadmium 112.4	indium 114.8	tin 118.7	antimony 121.8	tellurium 127.6	iodine 126.9	xenon 131.3
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ва	lanthanoids	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	T1	Pb	Bi	Po	At	Rn
caesium 132.9	barium 137.3		hafnium 178.5	tantalum 180.9	tungsten 183.8	rhenium 186.2	osmium 190.2	iridium 192.2	platinum 195.1	gold 197.0	mercury 200.6	thallium 204.4	lead 207.2	bismuth 209.0	polonium —	astatine -	radon —
87	88	89–103	104	105	106	107	108	109	110	111	112		114		116		
Fr	Ra	actinoids	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn		F1		Lv		
francium	radium		rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	copernicium		flerovium		livermorium		
_	_		_	_	_	_	_	_	_	-	_		_	L	_		

actinoids

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Но	Er	Tm	Yb	Lu
lanthanum 138.9	cerium 140.1	praseodymium 140.9	neodymium 144.4	promethium —	samarium 150.4	europium 152.0	gadolinium 157.3	terbium 158.9	dysprosium 162.5	holmium 164.9	erbium 167.3	thulium 168.9	ytterbium 173.1	lutetium 175.0
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
actinium -	thorium 232.0	protactinium 231.0	uranium 238.0	neptunium —	plutonium —	americium -	curium —	berkelium –	californium —	einsteinium –	fermium —	mendelevium -	nobelium —	lawrencium -