



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
 General Certificate of Education
 Advanced Subsidiary Level and Advanced Level

CANDIDATE
NAME

| |
|--|
| |
|--|

CENTRE
NUMBER

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

CANDIDATE
NUMBER

| | | | |
|--|--|--|--|
| | | | |
|--|--|--|--|



CHEMISTRY

9701/33

Advanced Practical Skills 1

May/June 2013

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
 Give details of the practical session and laboratory where appropriate, in the boxes provided.
 Write in dark blue or black pen.
 You may use a soft pencil for any diagrams, graphs or rough working.
 Do not use staples, paper clips, highlighters, glue or correction fluid.
DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.
 Electronic calculators may be used.
 You may lose marks if you do not show your working or if you do not use appropriate units.
 Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 12 and 13.
 A Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.
 The number of marks is given in brackets [] at the end of each question or part question.

| | |
|-------------------|--|
| Session | |
| | |
| Laboratory | |
| | |

| | |
|---------------------------|--|
| For Examiner's Use | |
| 1 | |
| 2 | |
| 3 | |
| Total | |

This document consists of **14** printed pages and **2** blank pages.

- 1 You are to determine the enthalpy change of the reaction between hydrochloric acid and sodium hydroxide by adding various volumes of acid and alkali and measuring the change in temperature.

FA 1 is $0.950 \text{ mol dm}^{-3}$ hydrochloric acid, HCl.

FA 2 is aqueous sodium hydroxide, NaOH.

(a) Method

- Support the plastic cup in a 250 cm^3 beaker.
- Using a measuring cylinder, transfer 25 cm^3 of **FA 1** into the cup and measure the temperature of the acid. Tilt the cup if necessary to cover the bulb of the thermometer.
- Record this initial temperature.

initial temperature of **FA 1** = °C

- Use a second measuring cylinder to transfer 10 cm^3 of **FA 2** and 25 cm^3 of water into a 100 cm^3 beaker.
- Add this mixture to the plastic cup and stir.
- Measure the maximum temperature reached and record this maximum temperature in the table below.
- Rinse out the plastic cup and shake it to remove excess water.
- Repeat the experiment, using the volumes of **FA 1**, **FA 2** and water shown in the table. Record the maximum temperature for each experiment.

| volume FA 1 / cm^3 | volume FA 2 / cm^3 | volume water / cm^3 | maximum temperature / °C |
|------------------------------------|------------------------------------|------------------------------|--------------------------|
| 25 | 10 | 25 | |
| 25 | 15 | 20 | |
| 25 | 20 | 15 | |
| 25 | 25 | 10 | |
| 25 | 30 | 5 | |
| 25 | 35 | 0 | |
| | | | |
| | | | |


You are going to plot a graph using these results to find the volume of **FA 2** that gives the greatest maximum temperature.

Before you plot the graph, choose two further volumes of **FA 2** that will allow you to find more precisely the volume that gives the greatest maximum temperature.

Record the volumes you choose, carry out the experiments and record the corresponding maximum temperatures, in the table. [2]

- (b) (i) On the grid below, plot the maximum temperature on the y -axis against the volume of **FA 2** on the x -axis.

For
Examiner's
Use



- (ii) Draw two straight lines of best fit on your graph, one to show where the temperature was increasing and the other after the greatest maximum temperature had been reached.
- (iii) Using your graph and the initial temperature recorded in (a), determine the maximum temperature **change** that could occur when 25 cm³ of **FA 1** react with **FA 2**.

maximum temperature **change** = °C
[5]

(c) Calculation

- (i) Calculate the energy needed to produce the temperature change in **(b)(iii)**.
(Assume that 4.3J of heat energy changes the temperature of 1.0 cm³ of solution by 1.0 °C.)

energy needed = J

- (ii) Calculate the number of moles of HCl used in each experiment.

moles of HCl = mol

- (iii) Calculate the enthalpy change, in kJ mol⁻¹, when 1 mole of HCl reacts with NaOH.

enthalpy change = kJ mol⁻¹
(sign) (value) [3]

[Total: 10]

- 2 The identity of a metal, M, can be found by titrating a solution of its carbonate with hydrochloric acid of known concentration.

For
Examiner's
Use

FA 3 is a solution of the metal carbonate, M_2CO_3 , of concentration 6.90 g dm^{-3} .

You are to dilute the hydrochloric acid that you used in **Question 1** and then titrate the carbonate solution with this acid.

(a) Method

Dilution of the acid

- Pipette 25.0 cm^3 of **FA 1** into the 250 cm^3 volumetric (graduated) flask labelled **FA 4**.
- Add distilled water to make the total volume 250 cm^3 .
- Stopper the flask and mix the contents thoroughly.

Titration

- Fill the burette with diluted hydrochloric acid, **FA 4**.
- Use a clean pipette to transfer 25.0 cm^3 of **FA 3** into a conical flask.
- Titrate **FA 3** with **FA 4** using the indicator provided.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is cm^3 .

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of **FA 4** added in each accurate titration.

| | |
|-----|--|
| I | |
| II | |
| III | |
| IV | |
| V | |
| VI | |
| VII | |


[7]

- (b) From your accurate titration results, obtain a suitable value to be used in your calculations. Show clearly how you obtained this value.

25.0 cm³ of **FA 3** required cm³ of **FA 4**
[1]

(c) **Calculation**

The equation for the reaction between hydrochloric acid and the metal carbonate is given below.



- (i) Calculate the number of moles of hydrochloric acid present in the volume in (b).

moles of HCl = mol

- (ii) Hence, calculate the number of moles of M₂CO₃ present in 25.0 cm³ of **FA 3**.

moles of M₂CO₃ = mol

- (iii) Calculate the concentration of M₂CO₃ in **FA 3** in mol dm⁻³.

concentration of M₂CO₃ = mol dm⁻³

- (iv) Use your answer to (iii), and the fact that **FA 3** contains 6.90 g dm⁻³, to determine the relative atomic mass, *A_r*, of M.

A_r of M =

- (v) Use your answer to (iv) and the Periodic Table on page 16 to suggest the identity of M.

M is [5]

(d) The concentration of a carbonate solution could be found using either the method in **Question 1** or that in **Question 2**.

(i) Suggest, and explain, which of the methods is more accurate.

.....
.....

(ii) For the method that you think is less accurate, suggest an improvement to the practical procedure that could be made to improve the accuracy.

.....
..... [2]

[Total: 15]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations.**

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

- (a) You are provided with a solid, **FA 5**. **FA 5** is a mixture that contains two anions and two cations.

To all your sample of **FA 5** in a boiling tube add 3 cm depth of distilled water. Shake the tube and filter the contents. Keep the solid residue for tests in (b) and the filtered solution for tests in (c).

- (b) (i) Open up the filter paper and scrape the residue into a boiling tube. Add dilute nitric acid, HNO_3 , using a dropping pipette until the solid **just** disappears. Record your observations and keep the solution for tests in (ii).

observations

.....

- (ii) Divide the solution from test (i) equally into **three** test-tubes.

To the first test-tube add aqueous sodium hydroxide, NaOH , until in excess. Record your observations.

observations

.....

Which cations, from those listed in the Qualitative Analysis Notes on page 12, would give these observations?

.....

- (iii) You are to devise tests that will positively identify which one of the cations you have suggested in (ii) is present. For each of the possible ions you should indicate the test and the expected result for each test in a suitable table in the space below.

Use the solutions in the second and third test-tubes to carry out these tests and record your observations in the space below.

Identify the cation present.

The cation present is

[7]

(c) To 1 cm depth of filtered solution from (a) in a test-tube add 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate. Record your observation.

observation

Which further reagent could be added to this test-tube to help you to confirm the nature of the anion present?

reagent

Carry out a test using this additional reagent. Record your observation and conclusion about the anion present.

observation

The anion present is [2]

(d) Using your observation in (b)(i) state which other anion is present in FA 5.

The anion present is [1]

- (e) Solutions **FA 6** and **FA 7** each contain one of the ions sulfite, SO_3^{2-} , sulfate, SO_4^{2-} , nitrite, NO_2^- , or nitrate, NO_3^- .

- (i) Carry out the tests in the table below to identify which ion is present in each solution.

| <i>test</i> | <i>observations</i> | |
|---|---------------------|-------------|
| | FA 6 | FA 7 |
| To 1 cm depth of solution in a boiling tube, add a small piece of aluminium foil and 1 cm depth of aqueous sodium hydroxide. Warm the mixture with care . | | |
| To 1 cm depth of solution in a test-tube, add a few drops of aqueous barium chloride or barium nitrate, then | | |
| add dilute hydrochloric acid. | | |
| To 1 cm depth of solution in a test-tube, add 1 cm depth of dilute hydrochloric acid. | | |

- (ii) From your observations, identify the anion present in each solution.

FA 6 contains

FA 7 contains

- (iii) What type of reaction takes place when a positive observation is seen with aluminium foil and aqueous sodium hydroxide in (i)?

.....
[5]

[Total: 15]

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

| ion | reaction with | |
|--|--|--|
| | NaOH(aq) | NH ₃ (aq) |
| aluminium, Al ³⁺ (aq) | white ppt. soluble in excess | white ppt. insoluble in excess |
| ammonium, NH ₄ ⁺ (aq) | no ppt. ammonia produced on heating | – |
| barium, Ba ²⁺ (aq) | no ppt. (if reagents are pure) | no ppt. |
| calcium, Ca ²⁺ (aq) | white ppt. with high [Ca ²⁺ (aq)] | no ppt. |
| chromium(III), Cr ³⁺ (aq) | grey-green ppt. soluble in excess giving dark green solution | grey-green ppt. insoluble in excess |
| copper(II), Cu ²⁺ (aq) | pale blue ppt. insoluble in excess | blue ppt. soluble in excess giving dark blue solution |
| iron(II), Fe ²⁺ (aq) | green ppt. turning brown on contact with air insoluble in excess | green ppt. turning brown on contact with air insoluble in excess |
| iron(III), Fe ³⁺ (aq) | red-brown ppt. insoluble in excess | red-brown ppt. insoluble in excess |
| lead(II), Pb ²⁺ (aq) | white ppt. soluble in excess | white ppt. insoluble in excess |
| magnesium, Mg ²⁺ (aq) | white ppt. insoluble in excess | white ppt. insoluble in excess |
| manganese(II), Mn ²⁺ (aq) | off-white ppt. rapidly turning brown on contact with air insoluble in excess | off-white ppt. rapidly turning brown on contact with air insoluble in excess |
| zinc, Zn ²⁺ (aq) | white ppt. soluble in excess | white ppt. soluble in excess |

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

2 Reactions of anions

| <i>ion</i> | <i>reaction</i> |
|---|---|
| carbonate, CO_3^{2-} | CO_2 liberated by dilute acids |
| chromate(VI), $\text{CrO}_4^{2-}(\text{aq})$ | yellow solution turns orange with $\text{H}^+(\text{aq})$; gives yellow ppt. with $\text{Ba}^{2+}(\text{aq})$; gives bright yellow ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| chloride, $\text{Cl}^-(\text{aq})$ | gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| bromide, $\text{Br}^-(\text{aq})$ | gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| iodide, $\text{I}^-(\text{aq})$ | gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$); gives yellow ppt. with $\text{Pb}^{2+}(\text{aq})$ |
| nitrate, $\text{NO}_3^-(\text{aq})$ | NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and <i>Al</i> foil |
| nitrite, $\text{NO}_2^-(\text{aq})$ | NH_3 liberated on heating with $\text{OH}^-(\text{aq})$ and <i>Al</i> foil; NO liberated by dilute acids (colourless $\text{NO} \rightarrow$ (pale) brown NO_2 in air) |
| sulfate, $\text{SO}_4^{2-}(\text{aq})$ | gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ or with $\text{Pb}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids) |
| sulfite, $\text{SO}_3^{2-}(\text{aq})$ | SO_2 liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids) |

3 Tests for gases

| <i>gas</i> | <i>test and test result</i> |
|-------------------------------|--|
| ammonia, NH_3 | turns damp red litmus paper blue |
| carbon dioxide, CO_2 | gives a white ppt. with limewater (ppt. dissolves with excess CO_2) |
| chlorine, Cl_2 | bleaches damp litmus paper |
| hydrogen, H_2 | “pops” with a lighted splint |
| oxygen, O_2 | relights a glowing splint |
| sulfur dioxide, SO_2 | turns acidified aqueous potassium dichromate(VI) from orange to green |

The Periodic Table of the Elements

| Group | | | | | | | | | | | | | | | | | | | | | |
|-------------------------------------|--------------------------------------|---------------------------------------|--------------------------------------|------------------------------------|---------------------------------------|--------------------------------------|-------------------------------------|-----------------------------------|-------------------------------------|-----------------------------------|-----------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|-------------------------------------|-------------------------------------|------------------------------------|--|--|--|---------------------------------|
| I | II | | | | | | | | | | | III | IV | V | VI | VII | 0 | | | | |
| | | | | | | | | | | 1.0 H Hydrogen 1 | | | | | | | | | | | 4.0 He Helium 2 |
| 6.9 Li Lithium 3 | 9.0 Be Beryllium 4 | | | | | | | | | | | 10.8 B Boron 5 | 12.0 C Carbon 6 | 14.0 N Nitrogen 7 | 16.0 O Oxygen 8 | 19.0 F Fluorine 9 | 20.2 Ne Neon 10 | | | | |
| 23.0 Na Sodium 11 | 24.3 Mg Magnesium 12 | | | | | | | | | | | 27.0 Al Aluminium 13 | 28.1 Si Silicon 14 | 31.0 P Phosphorus 15 | 32.1 S Sulfur 16 | 35.5 Cl Chlorine 17 | 39.9 Ar Argon 18 | | | | |
| 39.1 K Potassium 19 | 40.1 Ca Calcium 20 | 45.0 Sc Scandium 21 | 47.9 Ti Titanium 22 | 50.9 V Vanadium 23 | 52.0 Cr Chromium 24 | 54.9 Mn Manganese 25 | 55.8 Fe Iron 26 | 58.9 Co Cobalt 27 | 58.7 Ni Nickel 28 | 63.5 Cu Copper 29 | 65.4 Zn Zinc 30 | 69.7 Ga Gallium 31 | 72.6 Ge Germanium 32 | 74.9 As Arsenic 33 | 79.0 Se Selenium 34 | 79.9 Br Bromine 35 | 83.8 Kr Krypton 36 | | | | |
| 85.5 Rb Rubidium 37 | 87.6 Sr Strontium 38 | 88.9 Y Yttrium 39 | 91.2 Zr Zirconium 40 | 92.9 Nb Niobium 41 | 95.9 Mo Molybdenum 42 | Tc Technetium 43 | 101 Ru Ruthenium 44 | 103 Rh Rhodium 45 | 106 Pd Palladium 46 | 108 Ag Silver 47 | 112 Cd Cadmium 48 | 115 In Indium 49 | 119 Sn Tin 50 | 122 Sb Antimony 51 | 128 Te Tellurium 52 | 127 I Iodine 53 | 131 Xe Xenon 54 | | | | |
| 133 Cs Caesium 55 | 137 Ba Barium 56 | 139 La Lanthanum 57 * | 178 Hf Hafnium 72 | 181 Ta Tantalum 73 | 184 W Tungsten 74 | 186 Re Rhenium 75 | 190 Os Osmium 76 | 192 Ir Iridium 77 | 195 Pt Platinum 78 | 197 Au Gold 79 | 201 Hg Mercury 80 | 204 Tl Thallium 81 | 207 Pb Lead 82 | 209 Bi Bismuth 83 | Po Polonium 84 | At Astatine 85 | Rn Radon 86 | | | | |
| Fr Francium 87 | Ra Radium 88 | Ac Actinium 89 † | Rf Rutherfordium 104 | Db Dubnium 105 | Sg Seaborgium 106 | Bh Bohrium 107 | Hs Hassium 108 | Mt Meitnerium 109 | Uun Ununnilium 110 | Uuu Unununium 111 | Uub Ununbium 112 | | Uuq Ununquadium 114 | | Uuh Ununhexium 116 | | Uuo Ununoctium 118 | | | | |

*58-71 Lanthanides

†90-103 Actinides

Key

| |
|----------|
| a |
| X |
| b |

a = relative atomic mass
X = atomic symbol
 b = proton (atomic) number

| | | | | | | | | | | | | | |
|----------------------------------|--|-------------------------------------|-------------------------------|------------------------------------|------------------------------------|--------------------------------------|-----------------------------------|--------------------------------------|-----------------------------------|----------------------------------|-----------------------------------|-------------------------------------|------------------------------------|
| 140 Ce Cerium 58 | 141 Pr Praseodymium 59 | 144 Nd Neodymium 60 | Pm Promethium 61 | 150 Sm Samarium 62 | 152 Eu Europium 63 | 157 Gd Gadolinium 64 | 159 Tb Terbium 65 | 163 Dy Dysprosium 66 | 165 Ho Holmium 67 | 167 Er Erbium 68 | 169 Tm Thulium 69 | 173 Yb Ytterbium 70 | 175 Lu Lutetium 71 |
| Th Thorium 90 | Pa Protactinium 91 | U Uranium 92 | Np Neptunium 93 | Pu Plutonium 94 | Am Americium 95 | Cm Curium 96 | Bk Berkelium 97 | Cf Californium 98 | Es Einsteinium 99 | Fm Fermium 100 | Md Mendelevium 101 | No Nobelium 102 | Lr Lawrencium 103 |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.