UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2010 question paper for the guidance of teachers

9701 CHEMISTRY

9701/35

Paper 31 (Advanced Practical Skills), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	35

Question 1

Qu	estion	Sections	Indicative material	Mark	
1	(a)	MMO Collection	(i) Follows instructions with regard to weighings and mass of NaHCO ₃ 2 balance readings and mass of NaHCO ₃ and times and temperature readings 0–3 minutes at 1 minute intervals; 5–8 minutes at ½ minute intervals	1	
		PDO Recording	 (ii) All columns correctly labelled with appropriate unit shown. Must use solidus, brackets or describe unit fully in words. If units not included in column headings every entry must have the correct unit shown Accept min, mins or minutes 	1	
			(iii) Look at results here and in (d). All balance readings consistent to at least 1 decimal place. and All thermometer readings recorded to nearest 0.5°C. There must be at least one at 0.5 in (a).	1	[3]

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	35

Question	Sections	Indicative material	Mark	
(b)	PDO Layout	(i) Temperature of acid/solution in the beaker plotted on y-axis against time on x-axis. Clearly labelled axes (ignore units) [temp/time are minimum acceptable labels] but accept T I °C and t I min as labels. The unit is necessary in this case	1	
		 (ii) Uniform and sensible scales for candidate's choice of graph. Plotted points must be in at least 5 large squares on the temperature axis and 5 large squares on the time axis. 	1	
		 (iii) There should be a minimum of 5 plotted points between 5 and 8 minutes. Examiner then checks plotting of points at t_{0 min}, t_{5 min} and t_{8 min} and the plotting of any suspect point. If any of the t_{0 min}, t_{5 min} and t_{8 min} points is missing check the adjacent point. Points should be within ½ of a small square of the correct position and in the correct small square 	1	
	ACE Interpretation	(iv) Acceptable straight lines drawn – an acceptable straight line is one passing through the majority of points or has balanced points on either side of the line and correct values of the minimum and maximum temperatures at t = 4 minutes are read (to within ½ small square) from the graph. Extrapolation need not be drawn on the graph	1	[4]
(c)	ACE Interpretation	 (i) No mark. (ii) The candidate <u>correctly</u> calculates (to sig figs displayed) the moles of FA 1 used, (cand mass of FA 1/84) 	1	
	ACE	(iii) The candidate correctly divides their answer to (i) by their answer to (ii) and by 1000. Ignore errors in evaluation and sign.	1	
	Conclusions	(iv) Award this mark if the candidate has given a +ve sign and explains that: the reaction is endothermic or heat is absorbed in the reaction or the temperature falls during the reaction	1	[3]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	35

Question	Sections	Indicative material	Mark	
(d)	MMO Quality	Compare the two $\Delta T/m$ values (°C g ⁻¹) for the candidate's two experiments. Award three marks for a difference up to 0.2 Award two of these three marks for a difference of 0.2+ °C - 0.3 Award one of these three marks for a difference of	3	
		0.3+ °C $-$ 0.4 Compare the <u>standard</u> m/ ΔT value of 1.55°C g ⁻¹ with the closer value from the candidate's results. Award three marks for a difference up to 0.2 Award two of these three marks for a difference of 0.2+ °C $-$ 0.3 Award one of these three marks for a difference of 0.3+ °C $-$ 0.4	3	[6]
(e)	ACE Interpretation	 (i) No mark is awarded for this section but check that temperature rise from expt 1 has been used (ii) The candidate correctly calculates (to sig figs displayed) the moles of FA 2 used in expt 1. Do not award this mark if data from expt 2 has 	1	
		been used in (i) or (ii). (iii) The candidate correctly divides the answer to (e)(i) by the answer to (e)(ii) and by 1000. Ignore errors in evaluation and sign	1	
	ACE Conclusions	(iii) Award this mark if the candidate has given a –ve sign and explains that: the reaction is exothermic or heat is released in the reaction or	1	
	PDO Display	the temperature rises during the reaction Award this mark if working is shown in sections (c)(ii), (c)(iii), (e)(ii) and (e)(iii) Award this mark if the final answer in section (c)(iii) and	1	
		(e)(iii) is given to 2 or 3 sig fig.	ı	[5]
(f)	ACE Interpretation	Correctly calculates $\Delta H_{\rm decomposition}$ from candidate values in (c)(iii) and (e)(iii) . Allow (± 1) on final significant figure given The answer given must include a mathematically correct sign	1	[1]
(g)	ACE Interpretation	Correctly calculates the difference and the percentage error. Ignore significant figures.	1	[1]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	35

Question	Sections	Indicative material	Mark	
(h)	ACE Conclusions	Clearly described source of error (i) Heat loss / gain (ii) Precision of thermometer (iii) Acid spray (iv) Use of a glass beaker	1	
	ACE Improvements	 Specific improvement given with some attempt at justification. (i) Lid – prevents convection or evaporation Insulation – prevents conduction (ii) Use thermometer at 0.5°C or better, gives smaller % error. (iii) Lid – acts as a physical barrier (iv) Use of plastic cup – better insulator or lower (specific) heat capacity Do not credit either mark for answers referring to use of measurement of volume or measurement of mass. 	1	[2]
	Total			[25]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	35

Question 2

Ques	stion	Sections	Indicative material	Mark	
2 ((a)	MMO Decisions	(i) reagent 1 – chooses any specified acid to detect the carbonate present (name or formula may be in results table) and reagent 2 – chooses AgNO ₃ . Accept Ag+(aq) or soln containing Ag ⁺ as reagent. Also accept incorrect formulae for a compound, e.g. Ag(NO ₃) ₂ , providing the identity of the reagent is obvious.	1	
			 (ii) Explains significance of order in which reagents added. acid first – to remove carbonate from solution or after Ag⁺ – to dissolve any silver carbonate precipitated. Candidates must make clear the relationship of acid to silver carbonate. Do not award this mark if hydrochloric acid has been used in (i) In section (iii), assume reagents follow each other in the same test-tube unless otherwise stated. Allow lead(II) nitrate as the 2nd reagent providing it is used with nitric acid. 	1	
		MMO Collection	(iii) Addition of acid No reaction with FA 4, effervescence/bubbles/bubbling (or gas tested with limewater) for FA 5 and FA 6	1	
			Addition of Ag ⁺ (aq) Ignore any addition of NH ₃ (aq) after Ag ⁺ white ppt with FA 4, if added as first reagent or to a separate sample Allow off-white to brownish ppt with FA 6 or white ppt, insoluble in acid for FA 4, soluble in acid for FA 6 and insoluble/partially soluble in acid for FA 5 if added before the addition of acid or white ppt with FA 4 and FA 5 and no ppt with FA 6 if added after addition of acid Do not award this mark if hydrochloric acid has been used unless it has been stated that Ag ⁺ (aq) was added to a fresh sample Allow deductions from lead nitrate (as for silver salt) ONLY if nitric acid has been specified	1	

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2010	9701	35

Question	Sections	Indicative material	Mark	
2 (a) contd	ACE Conclusions	No ecf to be applied in these conclusions (iv) Give one mark for identifying carbonate in FA 5 and FA 6, with supporting evidence. Minimum acceptable evidence – gas with acid or off-white, white, cream. brownish ppt (silver carbonate) soluble in acid Do not award this mark from the colour of the precipitate alone. Give one mark for identifying chloride in FA 4 and FA 5 only, with supporting evidence. Minimum acceptable evidence – white ppt with Ag ⁺ (if insoluble in acid) but con if soluble. Do not award this mark from the colour of the precipitates with Ag ⁺ alone If no mark has been awarded in (iii) or (iv), allow one mark if evidence given is consistent with the ions	1	
		identified		[6]
FA 7 is A	!K(SO ₄) ₂ (aq); FA	8 is NiC <i>l</i> ₂ (aq); FA 9 is CrC <i>l</i> ₃ (aq); FA 10 is Pb(NO ₃) ₂ (aq)		
(b)	MMO Collection	For FA 7, records (i) white ppt, soluble in an excess of NaOH and white ppt, insoluble in excess ammonia	1	
		For FA 8, records (ii) green ppt, insoluble in an excess of NaOH and Dark or deep blue solution with (excess) ammonia or Initial blue solution, darkening with excess of the reagent	1	
		For FA 9 records (iii) grey-green ppt, soluble in an excess of NaOH to give a dark green solution	1	
		(iv) grey-green ppt, insoluble in excess ammonia	1	
		For FA 10, records (v) white ppt, soluble in an excess of NaOH and white ppt, insoluble in excess ammonia	1	

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper	
	GCE AS/A LEVEL – May/June 2010	9701	35	

(vi) Selects hydrochloric acid, sulfuric acid potassium iodide, potassium chromate (or dichromate) as suitable reagent Not barium chloride If no reagent is chosen, a retrospective De7 mark can be given here for testing gas evolved with limewater in (a) MMO Collection MMO Collection ACE Conclusions (vii) Observes appropriate ppt with chosen reagent and FA 10 but not with FA 7. Do not award this mark if reagent is added to any solution recorded as giving a coloured ppt or no ppt with NaOH or with NH3 or barium chloride is used No ecf to be applied in these conclusions Identifies all cations correctly: FA 7 is Al ³⁺ / aluminium FA 9 is CP ²⁺ / chromium(III) FA 10 is Pb ²⁺ / lead The correct ones must be identified for each solution. Gives appropriate supporting evidence for two of the three ions Allow evidence for reversed Al ²⁺ /Pb ²⁺ if barium chloride has been used Minimum evidence for each of the ions. Al ^{2+*} white ppt with NaOH and NH ₃ (aq); ppt soluble in excess NaOH, ppt insoluble in excess NaOH, and no reaction with HCl, H ₂ SO ₄ , KI Cr ^{2+*} (i) grey-green ppt with NaOH and NH ₃ (aq), insoluble in excess of the reagent, or (iii) grey-green ppt with NaOH and NH ₃ (aq), insoluble in excess of the reagent, or (iii) grey-green ppt with NaOH and NH ₃ (aq), insoluble in excess of the reagent, or (iii) Dark green solution with excess NaOH, ppt insoluble in excess NaOH, ppt insoluble in excess NaOH, and white ppt with NaOH and NH ₃ (aq); ppt soluble in excess NaOH, and white ppt with HCl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with Kl or H ₂ SO ₄ or yellow ppt with MaD or the proper in th	Question	Sections	Indicative material	Mark	
Collection FA 10 but not with FA 7. Do not award this mark if reagent is added to any solution recorded as giving a coloured ppt or no ppt with NaOH or with NH ₃ or barium chloride is used No ecf to be applied in these conclusions Identifies all cations correctly: FA 7 is A1 ²⁺ / aluminium FA 9 is Cr ²⁺ / chromium(III) FA 10 is Pb ²⁺ / lead The correct ions must be identified for each solution. Gives appropriate supporting evidence for two of the three ions Allow evidence for reversed A1 ³⁺ / Pb ²⁺ if barium chloride has been used Minimum evidence for each of the ions. A1 ³⁺ white ppt with NaOH and NH ₃ (aq); ppt isoluble in excess NaOH, ppt insoluble in excess of the reagent, or (ii) grey-green ppt with NaOH and NH ₃ (aq), insoluble in excess of the reagent, or (iii) grey-green ppt with NaOH, soluble in excess of the reagent, or (iii) grey-green ppt with NaOH, soluble in excess of the reagent, or (iii) Dark green solution with excess NaOH Pp2** white ppt with NaOH and NH ₃ (aq); ppt insoluble in excess NaOH, or yellow ppt with KI or H ₂ SO ₄ or yellow ppt with KI			iodide, potassium chromate (or dichromate) as suitable reagent Not barium chloride If no reagent is chosen, a retrospective De7 mark can be given here for testing gas evolved with limewater	1	
Identifies all cations correctly: FA 7 is A/3* / aluminium FA 9 is Cr3* / chromium(III) FA 10 is Pb2* / lead The correct ions must be identified for each solution. Gives appropriate supporting evidence for two of the three ions Allow evidence for reversed Al3*/Pb2* if barium chloride has been used Minimum evidence for each of the ions. Allow evidence for reversed Al3*/Pb2* if barium chloride has been used Minimum evidence for each of the ions. Allow evidence for two of th			FA 10 but not with FA 7. <u>Do not award this mark</u> if reagent is added to any solution recorded as giving a coloured ppt or no ppt	1	[7]
If an appropriate reagent has been chosen but no observations recorded in (b)(iii): Allow deduction of the identity of the ions if the observations have been correctly recorded in the evidence section Do not however award the mark for appropriate evidence from observations Allow evidence for reversed Al 3*/Pb 2* if barium chloride has been used	(c)	_	Identifies all cations correctly: FA 7 is Al ³⁺ / aluminium FA 9 is Cr ³⁺ / chromium(III) FA 10 is Pb ²⁺ / lead The correct ions must be identified for each solution. Gives appropriate supporting evidence for two of the three		
	reagent he chosen be observation (b)(iii): Allow decidentity of the observation been correcorded evidence Do not he the mark appropria	duction of the f the ions if vations have rectly in the section owever award for attentions to the section owever award for attention over a to the section of the section of the evidence —	Allow evidence for reversed Al ³⁺ /Pb ²⁺ if barium chloride has been used Minimum evidence for each of the ions. Al ³⁺ white ppt with NaOH and NH ₃ (aq); ppt soluble in excess NaOH, ppt insoluble in excess NH ₃ (aq) and no reaction with HCl, H ₂ SO ₄ , KI Cr ³⁺ (i) grey-green ppt with both NaOH and NH ₃ (aq), or (ii) grey-green ppt with NaOH, soluble in excess of the reagent, or (iii) grey-green ppt with NH ₃ (aq), insoluble in excess of the reagent, or (iii) Dark green solution with excess NaOH Pb ²⁺ white ppt with NaOH and NH ₃ (aq); ppt soluble in excess NaOH, ppt insoluble in excess NH ₃ (aq) and white ppt with HCl or H ₂ SO ₄ or		[2]
		Total	yellow ppt with KI		