



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

Candidates ans	swer on the Question Paper.		1 hour 15 minutes
Paper 3 (Exten	ided)		May/June 2015
CHEMISTRY			0620/31
CENTRE NUMBER		CANDIDATE NUMBER	
CANDIDATE NAME			

READ THESE INSTRUCTIONS FIRST

No Additional Materials are required.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



(a)	Coal is a solid fossil fuel.
	Name two other fossil fuels.
	[2]
(b)	Two of the elements present in a sample of coal are carbon and sulfur.
	A sample of coal was heated in the absence of air and the products included water, ammonia and hydrocarbons.
	Name three other elements present in this sample of coal.
	[2]
(c)	Sulfur, present in coal, is one major cause of acid rain. Sulfur burns to form sulfur dioxide which reacts with rain water to form sulfuric acid.
	(i) Describe how the high temperatures in vehicle engines are another cause of acid rain.
	[3]
	(ii) Give two harmful effects of acid rain.
	[2]
(d)	In 2010, a large coal-burning power station in the UK was converted to burn both coal and wood.
	Explain why the combustion of wood rather than coal can reduce the effect of the emissions from this power station on the level of carbon dioxide in the atmosphere.
	[3]
	[Total: 12]

2


	from the Blast Furnace is impure. It contains about 5% of impurities, mainly carbon, sulfur, n and phosphorus, which have to be removed when this iron is converted into steel.
(Explain how the addition of oxygen and calcium oxide removes these impurities. Include an equation for a reaction of oxygen and a word equation for a reaction of calcium oxide in this process.
	[5]
(b)	Mild steel is the most common form of steel. Mild steel contains a maximum of 0.3% of carbon.
I	High carbon steel contains 2% of carbon. It is less malleable and much harder than mild steel.
(i) Give a use of mild steel.
	[1]
(i	i) Suggest a use of high carbon steel.
	[1]
(ii	i) Explain why metals are malleable.
	[3]
(i	
	[2]
	[Total: 12]

3 (a) The reactions between metals and acids are redox reactions.


$$Zn + 2H^+ \rightarrow Zn^{2+} + H_2$$

(i) Which change in the above reaction is oxidation, Zn to Zn^{2+} or $2H^+$ to H_2 ? Give a reason for your choice.

- (ii) Which reactant in the above reaction is the oxidising agent? Give a reason for your choice.
- **(b)** The rate of reaction between a metal and an acid can be investigated using the apparatus shown below.



A piece of zinc foil was added to 50 cm³ of hydrochloric acid, of concentration 2.0 mol/dm³. The acid was in excess. The hydrogen evolved was collected in the gas syringe and its volume measured every minute. The results were plotted and labelled as graph 1.



The experiment was repeated to show that the reaction between zinc metal and hydrochloric acid is catalysed by copper. A small volume of aqueous copper(II) chloride was added to the acid before the zinc was added. The results of this experiment were plotted on the same grid and labelled as graph 2.

	(i)	Explain why the reaction mixture in the second experiment contains copper metal. Include an equation in your explanation.
		[2]
	(ii)	Explain how graph 2 shows that copper catalyses the reaction.
		[3]
(c)		ne first experiment was repeated using ethanoic acid, CH ₃ COOH, instead of hydrochloric d, how and why would the graph be different from graph 1?
		[4]
		[7]
(d)		culate the maximum mass of zinc which will react with 50cm^3 of hydrochloric acid, of centration 2.0mol/dm^3 .
		$Zn + 2HCl \rightarrow ZnCl_2 + H_2$
	Sho	ow your working.

[3]

[Total: 16]

The alc	cohols form a homologous series.	
(a) (i)	Give three characteristics which all members of a homologous series share.	
	[3	}]
(ii)	Give the name of the third member of this series.	
	name[1]
(iii)	Deduce the molecular formula of the alcohol whose $M_{\rm r}$ = 158. Show your working.	
	[2	2]
(b) Exp	plain why the following two alcohols are isomers.	
	$\begin{array}{c} CH_3 \\ CH_3 \\ CH_3 \\ CH_3 \\ CH_3 \end{array} \\ \begin{array}{c} CH_2 \\ CH_2 \\ CH_2 \\ CH_3 \\ \end{array} \\ \begin{array}{c} CH_2 \\ CH_2 \\ CH_3 \\ \end{array}$	

(c)	Thi	s question is based on typical reactions of butan-1-ol.
	(i)	When butan-1-ol, $CH_3-CH_2-CH_2-CH_2-OH$, is passed over the catalyst silicon(IV) oxide water is lost.
		Deduce the name and the structural formula of the organic product in this reaction.
		name
		structural formula
		[2
	(ii)	Suggest the name of the ester formed from butanol and ethanoic acid.
	()	[1
	(iii)	Butan-1-ol is oxidised by acidified potassium manganate(VII).
		Deduce the name and the structural formula of the organic product in this reaction.
		name
		structural formula
		re
		[2
		[Total: 13

- 5 The halogens are a group of non-metals in Group VII of the Periodic Table.
 - (a) The reactivity of the halogens decreases down the group.


Describe an experiment equation in your answer.									
The halogens form interha	alogen coi	mpounds	s. These a	are comp	ounds wh	nich coi	ntain tv	vo differ	ent

(b) The halogens form interhalogen compounds. These are compounds which contain two different halogens.

Deduce the formula of the compound which has the composition 0.013 moles of iodine atoms and 0.065 moles of fluorine atoms.

[2]

(c) Iodine reacts with chlorine to form a dark brown liquid, iodine monochloride.



$$I_2(s) + Cl_2(g) \rightarrow 2ICl(I)$$

When more chlorine is added and the tube is sealed, a reversible reaction occurs and the reaction comes to equilibrium.

$$ICl(I) + Cl_2(g) \rightleftharpoons ICl_3(s)$$
 dark brown yellow

(i)	Give another	example of a	reversible	reaction.
-----	--------------	--------------	------------	-----------

|--|

(ii) Explain the term equilibrium.

	[2]

(d)	Chlorine is removed from the tube and a new equilibrium is formed.
	Explain why there is less of the yellow solid and more dark brown liquid in the new equilibrium mixture.
	[2]
(e)	A sealed tube containing the equilibrium mixture is placed in ice-cold water. There is an increase in the amount of yellow solid in the equilibrium mixture.
	What can you deduce about the forward reaction in this equilibrium?
	$ICl(I) + Cl_2(g) \rightleftharpoons ICl_3(s)$
	Explain your deduction.
	[3]
	[Total: 13]

Aci	d-ba	se reactions are examples of proton transfer.
(a)	Eth	ylamine is a weak base and sodium hydroxide is a strong base.
	(i)	In terms of proton transfer, explain what is meant by the term weak base.
		[2]
	(ii)	Given aqueous solutions of both bases, describe how you could show that sodium hydroxide is the stronger base. How could you ensure a 'fair' comparison between the two solutions?
		[3]
(b)	Eth	ylamine reacts with acids to form salts.
		${\rm CH_3CH_2NH_2}$ + ${\rm HC}l$ \rightarrow ${\rm CH_3CH_2NH_3C}l$ ethylammonium chloride
	(i)	Complete the equation for the reaction between sulfuric acid and ethylamine. Name the salt formed.
		\dots CH ₃ CH ₂ NH ₂ + \dots \rightarrow \dots
		name of salt[3]
	(ii)	Amines and their salts have similar chemical properties to ammonia and ammonium salts
		Suggest a reagent that could be used to displace the weak base, ethylamine, from its sale ethylammonium chloride.
		[1]

c)	Gas	ses diffuse, which means that they move to occupy the total available volume.												
	(i)	Explain, using kinetic particle theory, why gases diffuse.												
		[2]												
	(ii)	When the colourless gases hydrogen bromide and ethylamine come into contact, a white solid is formed.												
		$CH_3CH_2NH_2(g) + HBr(g) \rightarrow CH_3CH_2NH_3Br(s)$ white solid												
		The following apparatus can be used to compare the rates of diffusion of the two gases ethylamine and hydrogen bromide.												
		gives off $CH_3CH_2NH_2(g)$ A B C C												
	CC	otton wool soaked in conc. hydrobromic acid												
		Predict at which position, A , B or C , the white solid will form. Explain your choice.												

.....[3]

[Total: 14]

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

DATA SHEET The Periodic Table of the Elements

Group																	
I	II		Group									III	IV	V	VI	VII	0
							1 H Hydrogen										4 He Helium 2
7 9 Li Be Lithium Beryllium 3 4												11 B Boron 5	12 C Carbon	14 N Nitrogen	16 O Oxygen 8	19 F Fluorine	20 Ne Neon
23 Na Sodium	24 Mg Magnesium 12											27 A 1 Aluminium 13	28 Si Silicon	31 P Phosphorus 15	32 S Sulfur	35.5 C1 Chlorine 17	40 Ar Argon
39 K Potassium 19	40 Ca Calcium 20	45 Sc Scandium 21	48 Ti Titanium 22	51 V Vanadium 23	52 Cr Chromium 24	55 Mn Manganese 25	56 Fe Iron	59 Co Cobalt 27	59 Ni Nickel	Cu Copper	65 Zn Zinc	70 Ga Gallium	73 Ge Germanium 32	75 As Arsenic	79 Se Selenium 34	Bromine 35	84 Kr Krypton 36
85 Rb Rubidium 37	88 Sr Strontium 38	89 Y Yttrium 39	91 Zr Zirconium 40	93 Nb Niobium 41	96 Mo Molybdenum 42	Tc Technetium 43	101 Ru Ruthenium 44	103 Rh Rhodium 45	106 Pd Palladium 46	108 Ag Silver	112 Cd Cadmium 48	115 In Indium 49	119 Sn Tin	122 Sb Antimony 51	128 Te Tellurium 52	127 I lodine 53	131 Xe Xenon 54
133 Cs Caesium 55	137 Ba Barium	139 La Lanthanum 57 *	178 Hf Hafnium 72	181 Ta Tantalum 73	184 W Tungsten 74	186 Re Rhenium 75	190 Os Osmium 76	192 Ir Iridium	195 Pt Platinum 78	197 Au Gold 79	201 Hg Mercury	204 T <i>l</i> Thallium 81	207 Pb Lead 82	209 Bi Bismuth	Po Polonium 84	At Astatine 85	Rn Radon 86
Fr Francium 87	226 Ra Radium	227 Ac Actinium 89 †															
*58-71 L	Actinoid	series		140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	Pm Promethium 61	150 Sm Samarium 62	152 Eu Europium 63	157 Gd Gadolinium 64	159 Tb Terbium 65	162 Dy Dysprosium 66	165 Ho Holmium 67	167 Er Erbium 68	169 Tm Thulium 69	173 Yb Ytterbium 70	175 Lu Lutetium 71
x = a a = relative atomic mass x = atomic symbol b = proton (atomic) number		232 Th Thorium 90	Pa Protactinium 91	238 U Uranium 92	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	Bk Berkelium 97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103		

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).