



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME							
CENTRE NUMBER					CANDIDATE NUMBER		
CHEMISTRY							0620/52
Paper 5 Practic	al Test					Мау	/June 2014
						1 hour	15 minutes
Candidates ans	swer on t	he Question	Paper.				
Additional Mate	erials:	As listed in	n the Con	fidential Instructions			

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
Total		

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 7 printed pages and 1 blank page.



1 You are going to investigate the temperature rise produced when different lengths of magnesium ribbon react with excess dilute sulfuric acid.

Read all the instructions below carefully before starting the experiments.

Instructions

You are going to carry out five experiments.

(a) Experiment 1

Using a measuring cylinder, pour 20 cm³ of dilute sulfuric acid into the beaker. Measure the initial temperature of the acid and record it in the table below. Add the 2 cm length of magnesium ribbon to the acid in the beaker, and stir the mixture with the thermometer.

Measure the highest temperature reached and record it in the table.

Remove the thermometer and rinse out the beaker with water.

(b) Experiment 2

Repeat Experiment 1, using the 3cm length of magnesium ribbon. Record the initial and highest temperatures in the table.

(c) Experiment 3

Repeat Experiment 1, using the 4 cm length of magnesium ribbon. Record the temperatures in the table.

(d) Experiment 4

Repeat Experiment 1, using the 6 cm length of magnesium ribbon. Record the temperatures in the table.


(e) Experiment 5

Repeat Experiment 1, using the 7 cm length of magnesium ribbon. Record the temperatures in the table.

Complete the table.

Experiment	initial temperature /°C	highest temperature /°C	change in temperature /°C
1			
2			
3			
4			
5			

(f) Plot the results you have obtained on the grid below. Draw a straight line of best fit through the points.



length of magnesium ribbon/cm

[4]

(g) From your graph, deduce the expected change in temperature if Experiment 1 was repeated using a 5 cm length of magnesium ribbon.

Show clearly on the graph how you worked out your answer.

[2	
.		4

(h) Give two observations when magnesium reacts with dilute sulfuric acid.

|--|

(i)	(i)	Which experiment gave the greatest change in temperature?
	(ii)	Suggest why the change in temperature was greatest in this experiment.
		[1]
(j)		at difference would be observed if Experiment 1 was repeated using an equal mass of gnesium powder? Explain your answer.
		[2]
(k)		lw a diagram of apparatus you could use to collect and measure the volume of gas given off ne reaction.
		[2]
(I)		te one source of error in the results obtained in Experiments 1-5. Give one improvement to uce this source of error.
	erro	or
	imp	provement[2]
		[Total: 21]

2 You are provided with mixture **E**. **E** consists of two solids, **F** and **G**. Solid **F** is water-soluble and solid **G** is insoluble.

Carry out the following tests on **E**, recording all of your observations in the table. Conclusions must **not** be written in the table.

	tests	observations
tests	on the mixture E	
(a)	Describe the appearance of the mixture.	[1]
\ · /	Place a little of mixture E in a test-tube.	
	Heat the mixture gently at first then more strongly for about 1 minute. After 1 minute, test the gas given off with damp pH indicator	
1	paper. Leave the test-tube to cool.	[3]
	Add the rest of mixture E to about 10 cm ³ of distilled water in a boiling tube. Stopper the boiling tube and shake the contents for about a minute. Filter the contents of the boiling tube. Keep the residue and the filtrate for the following tests.	
tests on the filtrate		
(c)	(i) To about 1 cm³ of the filtrate, add about 1 cm³ of aqueous sodium hydroxide. Gently heat the mixture. Test the gas given off with damp pH indicator paper.	[1]
(ii) Add about 1 cm ³ of silver nitrate solution to the second portion of the filtrate followed by about 1 cm ³ of dilute nitric acid.	[1]

		tests	observations
tes	ts or	n the residue	
(d)	res Us dilu	ing a spatula, transfer a little of the sidue from the filter paper to a test-tube. ing a teat pipette, add about 2 cm ³ of ute hydrochloric acid to the residue. Test e gas given off.	[2]
	the Sh	d an equal volume of distilled water to solution in the test-tube. ake the contents and divide into two rtions.	
(e)	(i)	Add several drops of aqueous sodium hydroxide to the first portion of the solution. Now add excess aqueous sodium hydroxide.	[3]
	(ii)	Add several drops of aqueous ammonia to the second portion. Now add excess aqueous ammonia.	[2]
(f)	Exp	lain your observations in test (b) .	
			[2]
(g)	Wha	at conclusions can you draw about solid F	
			[2]
(h)	Wha	at conclusions can you draw about solid G	3 ?
			[2]

[Total: 19]

BLANK PAGE

NOTES FOR USE IN QUALITATIVE ANALYSIS

Test for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I-) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH ₄ ⁺)	ammonia produced on warming	
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Test for gases

gas	test and test results	
ammonia (NH ₃)	turns damp red litmus paper blue	
carbon dioxide (CO ₂)	turns limewater milky	
chlorine (Cl ₂)	bleaches damp litmus paper	
hydrogen (H ₂)	'pops' with a lighted splint	
oxygen (O ₂)	relights a glowing splint	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.