

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

Candidates an	nswer on the Question Paper.		i iloui
Paper 6 Alternative to Practical			May/June 2013 1 hour
CHEMISTRY		062	
CENTRE NUMBER		CANDIDATE NUMBER	
CANDIDATE NAME			

READ THESE INSTRUCTIONS FIRST

No Additional Materials are required.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.



1 Electricity was passed through a solution of concentrated hydrochloric acid using the apparatus shown.


For Examiner's Use



(a)	Con	nplete the boxes to identify the parts of the apparatus labelled.	[2]
(b)	Des	scribe the test for hydrogen.	
	test		
	resu	ult	[2]
(c)		scribe how a sample of the gas given off at the positive electrode could be collected its volume measured.	ed
			[2]
(d)		experiment was repeated using a concentrated aqueous solution of sodium chloric ead of hydrochloric acid.	de
	(i)	State the name of the solution formed.	
			[1]
	(ii)	Give a test to show the presence of this product.	
			[1]
		[Total:	8]

- 2 A student found a recipe for making elderberry wine by fermentation.
 - 1 kg elderberries
 - 0.5 kg sugar
 - 10 g yeast granules
 - 3 dm³ water

The student decided to make some elderberry wine using the apparatus below.



The student carried out the following method.

- Step 1 The elderberries were crushed.
- The crushed elderberries and sugar were added to the water and the mixture was boiled for ten minutes. The crushed elderberries were then separated from the mixture.
- Step 3 Yeast was added to the liquid when it had cooled to room temperature.
- (a) Suggest the purpose of the airlock in the apparatus.

(b)	What apparatus could be used in Step 1?

......[1]

(c) Draw a labelled diagram of the apparatus used to separate the crushed elderberries from

the mixture in Step 2.

[2]

(d) Why was the yeast in Step 3 not added until the liquid was at room temperature?

(e)	(i)	State one observation during the fermentation.	E
		[1]	
	(ii)	Suggest how the rate of the fermentation reaction could be measured.	
		[2]	
(f)	Nar	me the method that could be used to separate ethanol from the fermented mixture.	
		[1]	
		[Total: 9]	

© UCLES 2013 0620/61/M/J/13

For Examiner's Use **3** A student investigated the reaction between two different solids, **C** and **D**, and excess dilute hydrochloric acid.

For Examiner's Use

Five experiments were carried out.

(a) Experiment 1

A measuring cylinder was used to pour 30 cm³ of dilute hydrochloric acid into a polystyrene cup. The temperature of the dilute hydrochloric acid was measured. 1 g of solid **C** was added to the dilute hydrochloric acid and the mixture stirred with a thermometer. The maximum temperature reached by the liquid mixture was measured.

(b) Experiment 2

The polystyrene cup was emptied and rinsed with water. Experiment 1 was repeated using 2 g of solid **C**.

(c) Experiments 3 and 4

Experiment 2 was repeated using 3 g and then 5 g of solid **C**.


Use the thermometer diagrams to record the results in the table below.

Complete the final column in the table.

experiment	mass of solid C	thermometer diagram	initial temperature of acid/°C	thermometer diagram	maximum temperature reached/°C	temperature difference /°C
1		30 -25 -20		30 -25 -20		
2		30 -25 -20		35 -30 -25		
3		30 -25 -20		35		
4		30 -25 -20		35 30 30 25		

[2]

Experiment 1 was repeated using solid \mathbf{D} . Use the thermometer diagrams to record the results in the spaces below.



initial temperature of dilute hydrochloric acid =°C


temperature change =°C

initial temperature of acid

final temperature of liquid mixture

final temperature of liquid mixture =°C

(e) Plot the results for Experiments 1, 2, 3 and 4 on the grid and draw a straight line graph.



[4]

(f)	(i)	From your graph, deduce the temperature of the solution when 6g of solid $\bf C$ is added to $30{\rm cm^3}$ of dilute hydrochloric acid. Show clearly on the grid how you worked out your answer.
		°C [2]
	(ii)	From your graph, deduce the mass of solid $\bf C$ that would give a temperature rise of 9 °C when added to 30 cm ³ of dilute hydrochloric acid.
		[2]
(g)	Wh	at type of chemical process occurs when solid D reacts with dilute hydrochloric acid?
		[1]
(h)	-	ggest the effect on the results if Experiment 3 was repeated using 60 cm³ of dilute rochloric acid.
		[2]
(i)		dict the temperature of the solution in Experiment 4 after 1 hour. Explain your answer.
		[2]
(j)	Wh	en carrying out the experiments, what would be one advantage and one disadvantage aking the temperature readings after exactly one minute?
	adv	antage
	disa	advantage
		[2]

[Total: 20]

For Examiner's Use **4** A mixture of two solids, **E** and **F**, was analysed.

Solid **E** was the water-soluble salt aluminium chloride, $AlCl_3$, and solid **F** was an insoluble salt.

The tests on the mixture and some of the observations are in the following table. Complete the observations in the table.

	tests	observations
boil The and	tilled water was added to the mixture in a ing tube. e contents of the boiling tube were shaken I filtered, keeping the filtrate and residue for following tests.	
test	s on the filtrate	
	e filtrate was divided into five portions in test-tubes.	
(a)	The first portion was used to describe the appearance of the filtrate.	appearance [1]
(b)	Several drops of aqueous sodium hydroxide were added to the second portion of the solution. Excess aqueous sodium hydroxide was then added to the test-tube.	[3]
(c)	Aqueous ammonia was added to the third portion, dropwise and then in excess.	[2]
(d)	To the fourth portion of the solution, dilute nitric acid and aqueous silver nitrate were added.	[2]
(e)	To the fifth portion of the solution, about 1 cm³ of dilute nitric acid and barium nitrate solution were added.	[1]


For Examiner's Use

tests	observations
tests on the residue	
(f) (i) To a little of the residue, dilute hydrochloric acid was added. The gas given off was tested.	rapid effervescence gas turned limewater milky
(ii) The residue was heated, gently then strongly.	solid changed colour from green to black

(g)	What conclusions can you draw about solid F ?
	[2

[Total: 11]

5 Identical pieces of steel were placed in two different boiling liquids for 12 hours. The graphs show how the mass of each piece of steel changed.



6	$\label{eq:copper} \begin{tabular}{l} Copper(II) oxide and carbon are both black solids. Copper(II) oxide reacts with dilute sulfuric acid to form aqueous copper(II) sulfate. Carbon does not react with dilute sulfuric acid. You are given a mixture of copper(II) oxide and carbon and access to dilute sulfuric acid. Plan an experiment to investigate the percentage of copper(II) oxide in the mixture. \\ \end{tabular}$	
	[6]	
	[Total: 6]	

For Examiner's Use

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.