

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

| | CANDIDATE NAME | | |
|-------|-------------------|----------------------------|---------------|
| | CENTRE NUMBER | CANDIDATE NUMBER | |
| * | | | |
| 0 3 | CHEMISTRY | | 0620/61 |
| 9 7 9 | Paper 6 Alterna | tive to Practical | May/June 2012 |
| 4 | | | 1 hour |
| 4 8 5 | Candidates ans | wer on the Question Paper. | |
| 1 4 4 | No Additional M | aterials are required. | |

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.


| For Examiner's Use | | |
|--------------------|--|--|
| 1 | | |
| 2 | | |
| 3 | | |
| 4 | | |
| 5 | | |
| 6 | | |
| Total | | |
| | | |

This document consists of **11** printed pages and **1** blank page.




1 A student reacted excess iron powder with sulfuric acid to prepare a solution of iron(II) sulfate.

The diagram shows the procedure followed in three stages.



2 Heat is given out when alcohols are burned.

A student used the apparatus below to find the amount of heat produced when four different alcohols, methanol, ethanol, propanol and butanol, were burned.



(a) Some methanol was put into the burner. The initial temperature of the water was measured. The burner was lit and allowed to burn for one minute. The flame was extinguished and the final temperature of the water was measured. The experiment was repeated with ethanol, propanol and butanol.

Use the thermometer diagrams to record the temperatures in the table on page 4. Complete the table by recording the temperature rise for each alcohol.

For

Examiner's Use

For Examiner's Use

| | 1 | | | | | |
|----------|----------------------------------|------------------------|----------------|------------------------|----------------|------------------------|
| | | initial | | final | | |
| alcohol | formula | thermometer diagram | temperature/°C | thermometer diagram | temperature/°C | temperature rise/°C |
| methanol | СН₃ОН | 25 20 | | 25 -20 | | |
| ethanol | C₂H₅OH | -25 -20 | | | | |
| propanol | C ₃ H ₇ OH | | | | | |
| butanol | C₄H₀OH | 25 20 | | | | |

[4]

5

(b) Plot the results obtained on the grid and draw a straight line graph.

50 40 30 temperature rise/°C 20 10 0 2 0 1 3 4 5 number of carbon atoms in the alcohol formula [4] (c) From your graph, work out the temperature rise expected if the experiment was repeated using pentanol, C₅H₁₁OH. Show clearly on the grid how you obtained your answer. [3] (d) Suggest the effect of using a copper can to contain the water instead of a boiling tube. Explain your answer. [Total: 13] For

Examiner's Use

© UCLES 2012

Coffee beans contain caffeine and other compounds. Caffeine is soluble in water and in

For

Examiner's

Use

3

trichloromethane, an organic solvent.

A student obtained crystals of caffeine by the following method.

BLANK PAGE

4 A student investigated the reaction between aqueous lead nitrate and aqueous potassium chloride.

For Examiner's Use

(a) One experiment was carried out.

Using a measuring cylinder, 3 cm³ of aqueous lead nitrate was poured into each of six test-tubes in a test-tube rack. The test-tubes were labelled A, B, C, D, E and F respectively.


A burette was filled with aqueous potassium chloride. A 1.0 cm³ sample of the aqueous potassium chloride was added to test-tube A.

A 2.0 cm³ sample of aqueous potassium chloride was added to test-tube B.

A 4.0 cm³, 5.0 cm³, 6.0 cm³ and 7.0 cm³ sample of aqueous potassium chloride was added to test-tubes C, D, E and F respectively.


Using a glass rod, the contents of the test-tubes were stirred. The contents of the test-tubes were left to stand for 10 minutes.

After 10 minutes, a ruler was used to measure the height of the solid in each test-tube. The diagrams show the six test-tubes in a rack. Use a ruler to measure the height of the solid in each test-tube in the diagram. Record the heights of the solid in the table.



| test-tube number | volume of aqueous potassium chloride / cm ³ | height of solid/mm |
|---------------------|--|--------------------|
| А | | |
| В | | |
| С | | |
| D | | |
| E | | |
| F | | |

[4]



(f) Predict what would happen if the experiment were continued using three further test-tubes with 8 cm³, 9 cm³ and 10 cm³ of aqueous potassium chloride. Explain your answer.
[2]
(g) What difference would be observed if the experiment was repeated using aqueous silver nitrate and aqueous potassium iodide?
[1]
(h) Explain one improvement the student could make to the experiment to obtain more accurate results.
improvement
explanation
[2]
[1]

For Examiner's Use

For Examiner's Use

Solid ${\boldsymbol W}$ was analysed. ${\boldsymbol W}$ was a carbonate salt. 5 The tests on solid **W**, and some of the observations, are in the following table. Complete the observations in the table.

Do not write any conclusions in the table.

| tests on solid W | |
|---|--|
| | |
| (a) Appearance of solid W. | vhite solid |
| | ied a white solid at the top he test-tube |
| The gas given off was tested with damp red litmus paper. litmus p | aper turned blue |
| (c) Dilute hydrochloric acid was added to solid W. | |
| The gas given off was tested. | |
| | [3] |
| (d) Dilute sodium hydroxide was added to solid W and the mixture heated. punge | ent gas given off |
| The gas given off was tested with damp pH pH indicator paper. | of gas = 10 |
| (e) Identify the gas given off in test (d). | |
| | [|
| (f) What conclusions can you draw about solid W? | |
| | [2 |
| | - |
| | [Total: 6 |

For Examiner's Use

STOP RUST!

Solutions of chemicals known as corrosion inhibitors are added to the water in steel radiators to reduce rust. You are provided with three different bottles of liquid corrosion inhibitors, **R**, **S** and **T**, and some steel nails. Plan an experiment to test if these inhibitors prevent the corrosion of steel and which of these inhibitors is the most effective.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

6