

MARK SCHEME for the May/June 2014 series

0606 ADDITIONAL MATHEMATICS

0606/23

Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.



Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0606	23

1	(i)	$500 = \frac{1}{2}r^2$ (1.6) 25 only	M1 A1	± 25 is A0
	(ii)	<i>their 25 + their 25 + their 25</i> $\times 1.6$ or better 90	M1 A1	<i>their 25</i> must be positive
2		$\log_x 3 = \frac{1}{\log_3 x}$ oe soi $u^2 - 4u - 12 = 0$ oe solve their 3 term quadratic in u Solve $\log_3 x = 6$ or $\log_3 x = -2$ oe 729 and $\frac{1}{9}$	B1 M1 M1 M1 A1	may be implied by $\log_x 3 = \frac{1}{u}$ oe condone sign errors
3	(i)	$\begin{pmatrix} 3 & 1 & 4 \\ 1 & 3 & 0 \end{pmatrix}$ and $\begin{pmatrix} 5 \\ 3 \\ 1 \end{pmatrix}$ or $(5 \ 3 \ 1)$ and $\begin{pmatrix} 3 & 1 \\ 1 & 4 \\ 4 & 0 \end{pmatrix}$ Multiplication of compatible matrices $\begin{pmatrix} 22 \\ 17 \end{pmatrix}$ or $(22 \ 17)$ as appropriate	B1 M1 A1	Must be correct shape from candidates product
	(ii)	$(1 \ 1)$ with $\begin{pmatrix} 22 \\ 17 \end{pmatrix}$ or $(22 \ 17)$ with $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	B1	

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0606	23

<p>4 (a) (i)</p> <p>(ii)</p> <p>(b) (i)</p> <p>(ii)</p> <p>(iii)</p>	  $50 \notin C$ $64 \in S \cap C$ $n(S') = 90$	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1ft</p> <p>B1</p>	<p>any Venn diagram showing three circles which do not all overlap</p> <p>ft only on use of $\not\subset$ and \subset instead of \notin and \in</p>
<p>5 (i)</p> <p>(ii)</p>	$(2\sqrt{2} + 4)^2 = 8 + 16\sqrt{2} + 16$ Correct completion Use $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ Multiply top and bottom by $2\sqrt{2} - 3$ $2 - \sqrt{2}$	<p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p>	$\left(\frac{(2\sqrt{2} + 4)}{2(2\sqrt{2} + 3)} \right)$ <p>Or $4\sqrt{2} - 6$</p>
<p>6</p>	Eliminate x or y Rearrange to quadratic in x or y $x^2 - 27x + 72 = 0$ or $y^2 + 9y - 90 = 0$ Factorise or solve 3 term quadratic $x = 3, x = 24$ or $y = 6, y = -15$ $y = 6, y = -15$ or $x = 3, x = 24$	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p>	

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0606	23

<p>7 (a)</p> $\frac{\frac{\sin \theta + \cos \theta}{\cos \theta} + \frac{\cos \theta + \sin \theta}{\sin \theta}}{\frac{1}{\cos \theta} + \frac{1}{\sin \theta}}$ <p>Clears the fractions in the numerator and denominator using common denominator</p> $\frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta + \cos \theta}$ <p>and completion</p> <p>(b) evidence of 13</p> $\sin x = \frac{5}{13}$ $\cos x = -\frac{12}{13}$		<p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>B1ft</p>	
<p>8 (i)</p> <p>Attempt to find $b^2 - 4ac$</p> <p>Completely correct argument</p> <p>(ii)</p> $m = 6(4) - 8(2) + 3$ $y - 10 = 11(x - 2) \text{ or } y = 11x - 12$ <p>(iii)</p> <p>Integrate to $2x^3 - 4x^2 + 3x(+c)$</p> $10 = 2(2)^3 - 4(2)^2 + 3(2) + c$ $y = 2x^3 - 4x^2 + 3x + 4 \text{ soi}$		<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B2,1,0</p> <p>M1</p> <p>A1</p>	<p>may be in formula or attempt to complete square</p> <p>dep on c being a genuine constant of integration</p> <p>ft on their 13</p>

Page 5	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0606	23

<p>9 (i)</p> <p>$(0, 7)$</p> <p>$m_{AB} = 2$</p> <p>perpendicular gradient = $-\frac{1}{2}$</p> <p>$y = -\frac{1}{2}x + 7$</p> <p>(ii)</p> <p>$m_{AB} = -1$</p> <p>$y = -x + 13$</p> <p>Solve their $y = -x + 13$ and $y = -\frac{1}{2}x + 7$</p> <p>$D(12,1)$</p> <p>Complete method for area</p> <p>84</p>		<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	
<p>10 (i)</p> <p>$\frac{d}{dx}(\sqrt{x^2 + 21}) = \frac{x}{\sqrt{x^2 + 21}}$</p> <p>Use of quotient rule</p> <p>$\frac{2\sqrt{(x^2 + 21)} - 2x \times \frac{x}{\sqrt{(x^2 + 21)}}}{(x^2 + 21)}$</p> <p>Multiply each term by $\sqrt{(x^2 + 21)}$</p> <p>$\frac{2(x^2 + 21) - 2x^2}{(x^2 + 21)^{\frac{3}{2}}}$ leading to $k = 42$</p> <p>(ii)</p> <p>$\frac{6}{k} \times \frac{2x}{\sqrt{x^2 + 21}}$</p> <p>Use limits in $C \times \frac{2x}{\sqrt{x^2 + 21}}$</p> <p>$\frac{8}{55}$ or 0.145</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>Alt method using product rule</p> <p>$\frac{d}{dx} \frac{1}{(\sqrt{x^2 + 21})^3} = \frac{-x}{(\sqrt{x^2 + 21})^3}$ is B1</p> <p>then M1 A1 as in quotient</p> <p>k must be a constant</p>	

Page 6	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0606	23

11	(i)	$\vec{OM} = \mathbf{a}$	B1	
		$\vec{MB} = 5\mathbf{b} - \mathbf{a}$	B1	
	(ii)	$\vec{ON} = 3\mathbf{b}$	B1	
		$\vec{AP} = \lambda(3\mathbf{b} - 2\mathbf{a})$	B1	
	(iii)	$\vec{MP} = \vec{MA} + \vec{AP}$	M1	
		$\mathbf{a} + \lambda(3\mathbf{b} - 2\mathbf{a})$	A1	
	(iv)	Put $\vec{MP} = \mu\vec{MB}$	M1	
		Equate components	M1	
		Solve simultaneous equations	M1	
		$\lambda = \frac{5}{7}$	A1	
12	(i)	$3 < f < 7$	B1,B1	If B0 then SC1 for $3 < f < 7$
	(ii)	$f(12) = 5$	B1	$f^2(x) = \sqrt{\sqrt{(x-3)+2}-3} + 2$ earns B1
		$(f(5) =) 2 + \sqrt{2}$	B1	
	(iii)	Clear indication of method $f^{-1}(x) = (x-2)^2 + 3$	M1 A1	condone $y = (x-2)^2 + 3$
	(iv)	$gf(x) = \frac{120}{\sqrt{(x-3)+2}}$	B1	
	Attempt to solve <i>their</i> $gf(x) = 20$	M1		
	$x = 19$	A1		