

**MARK SCHEME for the May/June 2013 series**

**0606 ADDITIONAL MATHEMATICS**

**0606/21**

Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – May/June 2013</b>	<b>0606</b>	<b>21</b>

## Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
  - A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
  - B Accuracy mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
  - The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
  - Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2, 1, 0 means that the candidate can earn anything from 0 to 2.

<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – May/June 2013</b>	<b>0606</b>	<b>21</b>

The following abbreviations may be used in a mark scheme or used on the scripts:

AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)

### Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.
OW –1, 2	This is deducted from A or B marks when essential working is omitted.
PA –1	This is deducted from A or B marks in the case of premature approximation.
S –1	Occasionally used for persistent slackness – usually discussed at a meeting.
EX –1	Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0606	21

<p><b>1</b></p>	$\frac{2 + 2 \sin^2 \theta}{\cos^2 \theta}$ $\frac{2}{\cos^2 \theta} = 2 \sec^2 \theta$ $\frac{\sin^2 \theta}{\cos^2 \theta} = 2 \tan^2 \theta$ <p><math>2 \sec^2 \theta = 2 + 2 \tan^2 \theta</math> and completion</p> <p><b>Or</b></p> $(\sec \theta + \tan \theta)^2 + (\sec \theta - \tan \theta)^2$ $2 \sec^2 \theta + 2 \tan^2 \theta$ $2(1 + \tan^2 \theta) + 2 \tan^2 \theta$ and completion <p><b>Or</b></p> $\frac{2 + 2 \sin^2 \theta}{\cos^2 \theta}$ $\frac{2(\sin^2 \theta + \cos^2 \theta) + 2 \sin^2 \theta}{\cos^2 \theta}$ $\frac{4 \sin^2 \theta}{\cos^2 \theta} = 4 \tan^2 \theta$ $\frac{2 \cos^2 \theta}{\cos^2 \theta} = 2$ and completion	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>[B1, B1]</b></p> <p><b>B1</b></p> <p><b>[B1]</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>[B1]</b></p>	<p>For all methods look for:</p> <ul style="list-style-type: none"> <li>– correct simplified expression</li> <li>– correct use of Pythagoras</li> <li>– use of <math>\tan = \frac{\sin}{\cos}</math></li> <li>– use of <math>\frac{1}{\cos} = \sec</math></li> </ul> <p>Award first 3 then last B1 for final expression from fully correct method.</p> <p>Inconsistent no angle used then –1 (can recover).</p> <p>If start from RHS award similarly.</p>
<p><b>2</b></p>	<p><b>(i)</b> 3.2</p> <p><b>(ii)</b> 15</p> <p><b>(iii)</b> uses area to find distance</p> <p>two of 40, 240 and 32</p> <p>312</p>	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>If split 2 or 3 correct formulae and must be attempting total area</p> <p>or <b>A2</b> for 312 from trapezium</p>

Page 5	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0606	21

3	$\frac{dy}{dx} = k \sin x \cos x$ $k = -8$ <p>Attempt to find <math>x</math> when <math>y = 8</math></p> $x = \frac{\pi}{4} \text{ (0.785)}$ <p>Uses <math>\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}</math></p> $-0.8 \text{ (not rounded)}$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>Must get to <math>x =</math> numerical value</p> <p><math>45^\circ = \mathbf{A0}</math> (but can still gain next 2 marks)</p> <p>Must use numerical value for <math>x</math> and 0.2 for <math>\frac{dx}{dt}</math></p> <p>(condone poor notation if correct terms multiplied)</p>
4	<p><b>(i)</b> Idea of modulus correct</p> <p><math>\frac{1}{2}</math> indicated on <math>x</math>-axis</p> <p>2 indicated on <math>y</math>-axis</p> <p><b>(ii)</b> <math>\frac{2}{3}</math> (0.667)</p> <p>Solve <math>4x - 2 = -x</math> or <math>(4x - 2)^2 = x^2</math></p> <p><math>\frac{2}{5}</math></p>	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>Two straight lines above and touching <math>x</math>-axis</p> <p>Must be a sketch</p> <p>Must be a sketch</p> <p>0.67 is <b>B0</b></p> <p>As far as <math>x =</math> numerical value</p> <p>SC: If drawn then <b>B1, B2</b> for exact answers only</p>
5	<p><b>(i)</b> <math>(QR = PS) = \frac{96 - 3x}{2}</math></p> <p>Area = <math>\left(\frac{96 - 3x}{2}\right) \times x</math></p> <p><b>(ii)</b> <math>\frac{dA}{dx} = \frac{96 - 6x}{2}</math> or <math>48 - 3x</math> o.e.</p> <p>Solving <math>\frac{dA}{dx} = \frac{96 - 6x}{2} = 0</math></p> <p><math>x = 16</math></p> <p><math>A = 384</math> and state maximum</p>	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>Can be implied by next statement</p> <p><b>AG</b></p> <p>As far as <math>x =</math> numerical value</p>

Page 6	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0606	21

6	<p>Applies quotient rule correctly</p> $\frac{(x-2)2x-(x^2+8)}{(x-2)^2}$ <p><math>y = 12</math></p> <p>Uses <math>m_1m_2 = -1</math></p> <p>(Gradient normal = <math>\frac{1}{2}</math>)</p> <p>Uses equation of line for <b>normal</b></p> $y-12 = \frac{1}{2}(x-4) \quad \text{or} \quad y = \frac{1}{2}x + 10$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>or product rule</p> $2x(x-2)^{-1} - (x^2+8)(x-2)^{-2}$ <p>If uses <math>y = mx + c</math> must find <math>c</math> for <b>M1</b></p>
7	<p>(i) <math>64 + 192x + 240x^2 + 160x^3</math> mark final answer</p> <p>(ii) Multiply out <math>(1 + 3x)(1 - x)</math></p> <p><math>1 + 2x - 3x^2</math> o.e.</p> <p><math>(1) \times (160) + (2) \times (240) + (-3) \times (192)</math> o.e.</p> <p>64</p> <p><b>Or</b></p> <p>Multiply out <math>(1 - x)(64 + 192x + 240x^2 + 160x^3)</math></p> <p>...<math>48x^2 - 80x^3</math>...o.e.</p> <p>Multiply by <math>1 + 3x</math></p> <p>64</p> <p><b>Or</b></p> <p><math>(1 + 3x)(64 + 192x + 240x^2 + 160x^3)</math></p> <p>...<math>816x^2 + 880x^3</math>...o.e.</p> <p>Multiply by <math>1 - x</math></p> <p>64</p>	<p><b>B3, 2, 1, 0</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>[M1]</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1]</b></p> <p><b>[M1]</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1]</b></p>	<p>3 terms correct earn <b>B2</b>; 2 terms correct earn <b>B1</b> Can be earned in (ii); <b>SC2</b> correct but unsimplified</p> <p>3 terms</p> <p>May be other variations: for first <b>M1</b> find <math>x^2</math> term or <math>x^3</math> term</p> <p>for second <b>M1</b> must produce all relevant terms</p>

Page 7	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0606	21

8	<p>Eliminates <math>y</math> (or <math>x</math>) and full attempt at expansion</p> $4x^2 - 8x - 96 = 0 \quad \text{or} \quad y^2 + 12y - 64 = 0$ <p>Factorise 3 term relevant quadratic</p> $x = -4 \text{ and } 6 \quad \text{or} \quad y = -16 \text{ and } 4$ $y = -16 \text{ and } 4 \quad \text{or} \quad x = -4 \text{ and } 6$ <p>Uses Pythagoras for relevant points</p> $22.4 \text{ or } \sqrt{500} \text{ or } 10\sqrt{5}$	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b>√</p> <p><b>M1</b></p> <p><b>A1</b></p>	<p>Or use correct formula</p> <p>cao</p>
9	<p><b>(i)</b> Attempt to solve 3 term quadratic</p> <p><math>-3</math> and <math>8</math></p> <p><math>-3 &lt; x &lt; 8</math></p> <p><b>(ii)</b> <math>4 &lt; x (&lt; 12)</math></p> <p><math>S \cup T = -3 &lt; x &lt; 12</math></p> <p><b>(iii)</b> <math>S \cap T = 4 &lt; x &lt; 8</math> or</p> <p><math>S' = -5 &lt; x \leq -3, 8 \leq x &lt; 12</math> and</p> <p><math>T' = -5 &lt; x \leq 4</math></p> <p><math>-5 &lt; x \leq 4</math></p> <p><math>8 \leq x &lt; 12</math></p>	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b>√</p> <p><b>B1</b>√</p>	<p>Condone <math>-3 &lt; x</math> AND <math>x &lt; 8</math></p> <p>Penalise confusion over <math>&lt;</math> and <math>\leq</math> (or <math>&gt;</math> and <math>\geq</math>) once only</p> <p><i>their 4</i></p> <p><i>their 8 (Ignore AND/OR etc.)</i></p>

Page 8	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0606	21

10	(i)	$\frac{\sin \alpha}{50} = \frac{\sin 58}{240}$ $\alpha = 10.2$ Bearing (0)21.8 or (0)22	M1 A1	Use of sin rule/cosine rule/resolving with 50, 240 and 58/32/122/148. Must be correct for A1
	(ii)	$V^2 = 240^2 + 50^2 - 2 \times 240 \times 50 \times \cos(122 - \alpha)$ $V = 263 \text{ awt}$ $T = \frac{500}{V}$ 114 or 1 hour 54 mins  <b>Or</b> $T = \frac{500 \cos 32}{240 \cos 21.8}$ $500 \cos 32$ $240 \cos 21.8$ 114 or 1 hour 54 mins	A1 A1√ M1 A1 M1 A1 [M1 B1 B1 A1]	√ for 32 – α  Correct use of sin rule/cosine rule/resolving  Can be in (i)  Only allow if V calculated from non right-angled triangle  Do not allow incorrect units  Alternative for part (ii) only Also can find distance for 240 (457) then 457/240
11	(i)	1	B1	Not a range for k, but condone x = 1 and x ≥ 1
	(ii)	f ≥ -5	B1	Not x, but condone y
	(iii)	Method of inverse  $1 + \sqrt{x+5}$	M1 A1	Do not reward poor algebra but allow slips  Must be f <sup>-1</sup> = ...or y = ...
	(iv)	f: Positive quadratic curve correct range and domain  f <sup>-1</sup> : Reflection of f in y = x	B1 B1√	Must cross x-axis  √their f(x) sketch Condone slight inaccuracies unless clear contradiction.
	(v)	Arrange f(x) = x or f <sup>-1</sup> (x) = x to 3 term quadratic = 0  4 only www	M1 A1	Allow x = 4 with no working. Condone (4, 4). Do not allow final A mark if -1 also given in answer



Page 9	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0606	21

12	(i)	$f(3) = (27 + 9 + 3a + b) = 0$ or $3a + b = -36$ $f(-1) = (-1 + 1 - a + b) = 20$ or $-a + b = 20$ Solve equations $a = -14, b = 6$	<b>M1</b> Equate $f(3)$ to 0  <b>M1</b> Equate $f(-1)$ to 20  <b>M1</b>  <b>A1</b> If uses $b = 6$ then <b>M0, A0</b> Need both values for <b>A1</b>
	(ii)	Find quadratic factor  $x^2 - 4x - 2$  Use quadratic formula or completing square on relevant 3 term quadratic  $\frac{-4 \pm \sqrt{16 + 8}}{2}$ or better  $-2 \pm \sqrt{6}$ isw	<b>M1</b> If division, must be complete with first 2 terms correct If writes down, must be $(x^2 + kx - 2)$  <b>A1</b>  <b>M1</b> If completing square, must reach $\left(x + \frac{k}{2}\right)^2 = 2 \pm \left(\frac{k}{2}\right)^2$  <b>A1√</b>  <b>A1</b> cao