

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME				
 CENTRE NUMBER			CANDIDATE NUMBER	
	MATHEN	IATICS		0606/22
Paper 2				May/June 2012
				2 hours
Candidates ans	swer on t	he Question Paper.		

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 80.

For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
Total		

This document consists of 16 printed pages.



UNIVERSITY of CAMBRIDGE International Examinations

[Turn over

### Mathematical Formulae

#### 1. ALGEBRA

Quadratic Equation

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

**Binomial Theorem** 

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$
  
where *n* is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!}$ 

### 2. TRIGONOMETRY

Identities

$$sin2 A + cos2 A = 1$$
$$sec2 A = 1 + tan2 A$$
$$cosec2 A = 1 + cot2 A$$

Formulae for  $\Delta ABC$ 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2} bc \sin A$$

		5		
1	It is given that $P$ is the set of prime numbers, $S$ is the set of square numbers and $N$ is the set numbers between 10 and 90. Write each of the following statements using set notation.			
	(i)	7 is a prime number.	[1]	
	(::)	e is not a course number	F13	
	(ii)	8 is not a square number.	[1]	
	(iii)	There are 6 square numbers between 10 and 90.	[1]	
2	(i)	Given that $y = \sqrt{(4x+1)^3}$ , find $\frac{dy}{dx}$ .	[2]	

(ii) Hence find the approximate increase in y as x increases from 6 to 6 + p, where p is small. [2]

3 Find the values of *m* for which the line y = mx - 5 is a tangent to the curve  $y = x^2 + 3x + 4$ .

4

4 In a competition the contestants search for hidden targets which are classed as difficult, medium or easy. In the first round, finding a difficult target scores 5 points, a medium target 3 points and an easy target 1 point. The number of targets found by the two contestants, Claire and Denise, are shown in the table.

For Examiner's Use

Target	Difficult	Medium	Easy
Claire	4	1	7
Denise	2	5	1

In the second round, finding a difficult target scores 8 points, a medium target 4 points and an easy target 2 points. In the second round Claire finds 2 difficult, 5 medium and 2 easy targets whilst Denise finds 4 difficult, 3 medium and 6 easy targets.

(i) Write down the sum of two matrix products which, on evaluation, would give the total score for each contestant. [3]

(ii) Use matrix multiplication and addition to calculate the total score for each contestant. [2]

- 5 It is given that x 2 is a factor of  $f(x) = x^3 + kx^2 8x 8$ .
  - (i) Find the value of the integer *k*.

(ii) Using your value of k, find the non-integer roots of the equation f(x) = 0 in the form  $a \pm \sqrt{b}$ , where a and b are integers. [5]

For Examiner's Use

[2]

6	(a)	Find the coefficient of $x^3$ in the expansion of	For Examiner's
		(i) $(1-2x)^7$ , [2]	Use

(ii) 
$$(3+4x)(1-2x)^7$$
. [3]

(**b**) Find the term independent of x in the expansion of 
$$\left(x + \frac{3}{x^2}\right)^6$$
. [3]

7 The table shows experimental values of variables *x* and *y*.

X	5	30	150	400
у	8.9	21.9	48.9	80.6

(i) By plotting a suitable straight line graph, show that y and x are related by the equation  $y = ax^b$ , where a and b are constants. [4]

			+++++++++++++++++++++++++++++++++++++++	
		+++++++++++++++++++++++++++++++++++++++	+++++++++++++++++++++++++++++++++++++++	
		+++++++++++++++++++++++++++++++++++++++	+++++++++++++++++++++++++++++++++++++++	
		<del>· · · · · · · · · · · · · · · · · · · </del>	****	┠┼┼┼┼┼┼┼┼╂┼┼┼┼┼┼┼┤
	┝╊┾┼┼┼┼┼┼┼╂┼┼┼┤	+++++++++++++++++++++++++++++++++++++++	****	
	┝╋┿┽┿┿┿┿┿┿┿╋┿┿┿┥	+++++++++++++++++++++++++++++++++++++++	+++++++++++++++++++++++++++++++++++++++	+ + + + + + + + + + + + + + + + + + +
	┝╉┼┼┼┼┼┼┼┼╂┼┼┼┤	<del></del>	+++++++++++++++++++++++++++++++++++++++	┠┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┤
	┝╋┾┾┾┾┾┾┾┾╋┾┾┾┥	<del>· · · · · · · · · · · · · · · · · · · </del>	****	┠┼┼┼┼┼┼┼╂┼┼┼┼┼┼┼┤
	┝┲┼┼┼┼┼┼┼┼┼┼┼┼	<del>·····································</del>		
		+++++++++++++++++++++++++++++++++++++++	+++++++++++++++++++++++++++++++++++++++	
	┝╉┼┼┼┼┼┼┼┼╂┼┼┼┤	<del>· · · · · · · · · · · · · · · · · · · </del>	┽┼┽┼┼┼╂┼┼┼┼┼┼┼┼	┠┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼
	┝╉┼┿┼┼┼┼┼┼╂┼┼┼┤	<del>╶╎╎╎╎╎╏╎╎╎╎╎╎╎╎╎╎</del>	****	┠┼┼┼┼┼┼┼╂┼┼┼┼┼┼┼┤
	┝╉┼┼┼┼┼┼┼┼╂┼┼┼┤	┼┼┼┼╂┼┼┼┼┼┼┼╂┼┼	<del>· · · · · · · · · · · · · · · · · · · </del>	
		*****	****	
		+++++++++++++++++++++++++++++++++++++++	****	
		<del></del>	****	
			+++++++++++++++++++++++++++++++++++++++	
		+++++++++++++++++++++++++++++++++++++++	****	
		*****		
+ + + + + + + + + + + + + + + + + + +	┝╋┼┼┼┼┼┼┼┼╂┼┼┼┤	<del>· · · · · · · · · · · · · · · · · · · </del>	+++++++++++++++++++++++++++++++++++++++	+ + + + + + + + + + + + + + + + + + +
	┝╉┼┼┼┼┼┼┼┼┼┼┼┼	<del>· · · · · · · · · · · · · · · · · · · </del>	<del>· · · · · · · · · · · · · · · · · · · </del>	
	┝╉┽┼┼┼┼┼┼┼╂┼┼┼┤	+++++++++++++++++++++++++++++++++++++++	+++++++++++++++++++++++++++++++++++++++	+ + + + + + + + + + + + + + + + + + +
	┝╉┼┼┼┼┼┼┼┼╂┼┼┼┤	<del>· · · · · · · · · · · · · · · · · · · </del>	<del>· · · · · · · · · · · · · · · · · · · </del>	┠┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼┤
	┝╋┼┼┼┼┼┼┼┼╂┼┼┼┤	*****	****	
		<del>╶╷╷╷╷╏╎╷╷╷╷╷╷╷╷╷</del>	┽┼┼┼┼┼╂┼┼┼┼┼┼┼┼	
	┝╋┽┼┼┼┼┼┼┼╂┼┼┼┤			+ + + + + + + + + + + + + + + + + + +
	┝╉┼┼┼┼┼┼┼┼╂┼┼┼┤	<del>· · · · · · · · · · · · · · · · · · · </del>	+++++++++++++++++++++++++++++++++++++++	+ + + + + + + + + + + + + + + + + + +
	┝╋┾┼┼┼┼┼┼┼╂┼┼┼┤	<del>· · · · · · · · · · · · · · · · · · · </del>	<del>· · · · · · · · · · · · · · · · · · · </del>	┠┼┼┼┼┼┼┼╂┼┼┼┼┼┼┼┤
		+++++++++++++++++++++++++++++++++++++++		
+ + + + + + + + + + + + + + + + + + +	┝╉┽┼┼┼┼┼┼┼╂┼┼┼┤	+++++++++++++++++++++++++++++++++++++++	+++++++++++++++++++++++++++++++++++++++	++++++++++++++++++++++++++++++++++++
	┝╉┼┼┼┼┼┼┼┼╂┼┼┼┤	<del>· · · · · · · · · · · · · · · · · · · </del>	+++++++++++++++++++++++++++++++++++++++	+ + + + + + + + + + + + + + + + + + +
	┝╋┼┼┼┼┼┼┼┼╂┼┼┼┤	*****	****	
+ + + + + + + + + + + + + + + + + + +	┝╋┽┽┽┽┽┽┼┼┼╋┼┼┼┤	<del></del>	+++++++++++++++++++++++++++++++++++++++	┠┼┼┼┼┼┼┼┼┼┼┼┼┼┼┼

For Examiner's Use (ii) Use your graph to estimate the value of a and of b.

(iii) Estimate the value of y when x = 100.

[2]

[4]

For Examiner's Use 8 An open rectangular cardboard box with a square base is to have a volume of 256 cm<sup>3</sup>. Find the dimensions of the box if the area of cardboard used is as small as possible. [7]

For Examiner's Use

9 (a) Solve the equation	
--------------------------	--

(i)  $3\sin x - 5\cos x = 0$  for  $0^\circ < x < 360^\circ$ ,

(ii)  $5\sin^2 y + 9\cos y - 3 = 0$  for  $0^\circ < y < 360^\circ$ .

[5]

For Examiner's

Use

[3]

 $\begin{bmatrix} 4 \end{bmatrix} \xrightarrow{F} Exan$ 

10 (a) A team of 7 people is to be chosen from 5 women and 7 men. Calculate the number of different ways in which this can be done if

12

- (i) there are no restrictions,
- (ii) the team is to contain more women than men.

[1]

(b) (i) How many different 4-digit numbers, less than 5000, can be formed using 4 of the 6 digits 1, 2, 3, 4, 5 and 6 if no digit can be used more than once? [2]

For Examiner's Use

(ii) How many of these 4-digit numbers are divisible by 5?

[2]

11 Answer only **one** of the following two alternatives.

### EITHER



The diagram shows part of the curve  $y = \sin \frac{1}{2}x$ . The tangent to the curve at the point  $P\left(\frac{3\pi}{2}, \frac{\sqrt{2}}{2}\right)$  cuts the *x*-axis at the point *Q*.

(i) Find the coordinates of Q.

[4]

For Examiner's Use

(ii) Find the area of the shaded region bounded by the curve, the tangent and the *x*-axis. [7]

#### OR

(i) Given that 
$$y = xe^{-x}$$
, find  $\frac{dy}{dx}$  and hence show that  $\int xe^{-x} dx = -xe^{-x} - e^{-x} + c.$  [4]



The diagram shows part of the curve  $y = xe^{-x}$  and the tangent to the curve at the point  $R\left(2, \frac{2}{e^2}\right)$ . (ii) Find the area of the shaded region bounded by the curve, the tangent and the *y*-axis. [7]

Start your answer to Question 11 here.				For
EITHER			Examiner's Use	
Indicate which question you are answering.	OR		-	
	UK			
		•••••		
		••••••		
		•••••		
		•••••		
		•••••		
		••••••		
		•••••		
		•••••		
		•••••		

Continue your answer here if necessary.	For
	Examiner's Use

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

### www.theallpapers.com

16