

Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

BIOLOGY 9700/52

Paper 5 Planning, Analysis and Evaluation

May/June 2017

MARK SCHEME
Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

 ${\rm \rlap{R}\hskip-1pt B}$ IGCSE is a registered trademark.



Mark scheme abbreviations

; separates marking points

I alternative answers for the same point

R reject

A accept (for answers correctly cued by the question, or extra guidance)

AW alternative wording (where responses vary more than usual)

underline actual word given must be used by candidate (grammatical variants accepted)

max indicates the maximum number of marks that can be given

ora or reverse argument ecf error carried forward

I ignore

mp marking point (with relevant number)

© UCLES 2017 Page 2 of 8

| Question | Answer | Marks | Guidance |
|-----------|---|-------|--|
| 1(a) | independent variable: concentration of caffeine; dependent variable: number of (heart) beats per unit time / heart rate; | | |
| 1(b)(i) | use two tablets; add 1 dm³ or 1000 ml/cm³ or 1 litre (distilled) water; | | A any correct proportions of water and tablets whatever the total volume, e.g. one tablet in 500 cm ³ / 0.5 dm ³ water |
| 1(b)(ii) | to keep it in one position / to stop it from moving / swimming (to make it easier to count the heart beat); | 1 | |
| 1(b)(iii) | max 6 of: ref. to a method of diluting 100 mg dm⁻³ caffeine solution, e.g. proportional / simple / serial dilution or a description and minimum of 4 additional dilutions; ref. to concentrations from 100 mg dm⁻³ downwards with correct | 6 | proportional / simple: (100), 80, 60, 40, 20, |
| | units; values stated <u>must</u> correspond to the dilution method chosen | | (0) mg dm ⁻³ ser <u>ial</u> : (100), 50, 25, 12.5, 6.25 / (100), 10, 1, 0.1, 0.01 mg dm ⁻³ must have a minimum of 3 others between 0.0 and 100.0 mg dm ⁻³ |
| | 3 use of water / 0 mg dm ⁻³ as a <u>control</u> ; | | |
| | 4 allow <i>Daphnia</i> to acclimatise after adding caffeine / to absorb the caffeine; | | |
| | 5 ref. to method of counting number of heart beats, e.g. clicker counter / tally counter / record dots on paper and count / video; | | |
| | 6 use of same period of time (for counting; | | standardising variables (mp6–mp8) – must be clear that all the concentrations have been tested |
| | same volume / same number of drops of caffeine solution added to each slide; if a value stated must be max 1 cm³ or 5 drops | | or one concentration has been tested more than once on <u>Daphnia</u> |

© UCLES 2017 Page 3 of 8

| Question | Answer | | | Guidance |
|----------|---|---|---|--|
| | 8 use the same organism / same size / same length / same age / same species / same type <i>Daphnia</i> for all caffeine concentrations; 9 ref. to a minimum of three replicates and calculate a mean or identify / eliminate / remove / ignore anomalies or outliers; 10 description of ethical treatment of live <i>Daphnia</i> AW, e.g. careful handling (when being moved) to minimise damage / return to tank promptly after testing / minimum time in caffeine solution; | | | |
| | | | | |
| | | | | |
| | 11 low risk experiment/suitable is allergy to caffeine and gloves; | nazard and safety precaution, e.g. | | |
| 1(b)(iv) | source of error is max 1 and must be clearly stated improvement is max 1 and must match the source of error | | 2 | A any other valid source of error and a suitable improvement I ref. to magnification used |
| | error | improvement | | |
| | heat from light in microscope; | turn lamp on only when needed / heat shield; | | |
| | evaporation of water from slide; | use a cover slip / top up with same solution; | | |
| | animals are stressed; | handle only when needed / minimise time in experimental conditions; | | |
| | cumulative effect of caffeine (on one <i>Daphnia</i>); | allow recovery time / use different Daphnia; | | |
| | difficulty in counting; | any suitable improvement, e.g. video and slow down; | | |
| | no time allowed for caffeine absorption; | have a time delay before counting; | | |
| | drop size varies ; | use a known volume of caffeine solution; | | |

© UCLES 2017 Page 4 of 8

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| 1(c) | Daphnia belong to a different phylum / data collected was not from humans; | 1 | A any ref. to differences in heart structure of humans and Daphnia |
| 1(d)(i) | (2.4 mg 100 cm ⁻³ cola, trial 3) <u>228</u> ; | 1 | |
| 1(d)(ii) | max 2 of: range of concentration too narrow; | 2 | |
| | no data for caffeine at 0.0 / below 2.4 / above 6.0 mg cm ⁻³ ; | | |
| | not enough concentration / only 4 concentrations; | | |
| | there is overlap between some of data collected for 4.8 and 6.0 mg cm ⁻³ ; | | |
| | idea that proportional increases in concentration should give a proportional increase in heart rate; | | |

© UCLES 2017 Page 5 of 8

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| 2(a) | there is no <u>significant</u> correlation / relationship / association between the percentage / proportion of cyanogenic <i>T. repens</i> and (increasing mean January) temperature ; | 1 | |
| 2(b)(i) | column 3 completed correctly; | 2 | ecf for column 6 from errors in column 3 |
| | column 6 completed correctly; | | |

| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|-------------|---|---|---------------------------------------|--|--------------------------|-------|
| location | percentage of cyanogenic <i>T.repens</i> plants | rank of percentage of cyanogenic <i>T.repens</i> plants | mean January temperature /°C | rank of mean January temperature | difference in rank, D | D^2 |
| Almora | 85 | 8 | 12.2 | 8 | 0 | 0 |
| Fairbanks | 5 | 2 | -23.9 | 1 | 1 | 1 |
| Karaj River | 64 | 5 | 4.4 | 6 | -1 | 1 |
| Konosu | 50 | 4 | 4.2 | 5 | -1 | 1 |
| Lennoxville | 71 | 7 | -10.0 | 4 | 3 | 9 |
| Mandan | 33 | 3 | -12.8 | 3 | 0 | 0 |
| Novosibirsk | 0 | 1 | -19.4 | 2 | -1 | 1 |
| Pretoria | 68 | 6 | 10.0 | 7 | -1 | 1 |
| Rabat | 100 | 9; | 12.5 | 9 | 0; | 0 |
| | | | | | $\Sigma D^2 =$ | 14 |

© UCLES 2017 Page 6 of 8

| Question | Answer | Marks | Guidance |
|-----------|---|-------|--|
| 2(b)(ii) | $r_s - 1 - \frac{(6 \times 14)}{(9^3 - 9)}$; | 2 | max 1 if correct answer is given to more than 2 d.p. |
| | $r_s - 1 - \frac{(84)}{(720)}$ | | |
| | $r_{\rm s} = 0.88$; | | |
| 2(b)(iii) | calculated value / 0.88 , is greater than, the critical value / 0.68 or critical value / 0.68 , is less than, the calculated value / 0.88 ; | 1 | ecf from incorrect answer in 2(b)(ii) |
| 2(b)(iv) | max 1 of: idea that cyanogenic plants grow better at higher temperature ; | 1 | must be comparative |
| | idea that cyanogenic plants more able to survive grazing (by herbivores); | | |
| | idea that cyanogenic plants produce more hydrogen cyanide which, reduces grazing / kills (more), herbivores ; | | |

© UCLES 2017 Page 7 of 8

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| 3(a) | max 3 of: same location / area used; | 3 | I species of vole |
| | same time of year / same two weeks in August; | | |
| | traps were equally spaced (along the transect); | | |
| | along same transects / transects were at the same places; | | |
| | numbers calculated per 1000 traps / same number of traps were used; | | |
| 3(b) | 1 $q^2 = 0.16 \text{ or } \frac{8}{50} \text{ or } \frac{4}{25} \text{ or } 16\%$ | 3 | max 2 if answer not rounded or p is incorrect A answers in equation as percentages |
| | OR $q = 0.4$ or $\frac{2}{5}$ or 4%; | | |
| | derives 2pq correctly from a clearly stated value of p and a clearly stated value of q; | | 2 ecf if <i>q</i> is incorrect (e.g. <i>q</i> = 0.16) but then correctly used to get <i>2pq</i> |
| | 3 in 1997 heterozygous voles = (0.48×60) = 29 voles; | | 3 ecf (any number) × 60 (from graph) <u>and</u> a <u>whole</u> number rounded correctly |

© UCLES 2017 Page 8 of 8