Cambridge International AS & A Level

Cambridge International Advanced Subsidiary and Advanced Level

	CANDIDATE NAME												
	CENTRE CANI NUMBER NUM	DIDATE BER											
* 5	CHEMISTRY	9701/33											
8 5	Paper 3 Advanced Practical Skills 1	February/March 2017											
⁸		2 hours											
	Candidates answer on the Question Paper.												
÷ 5 2 4	Additional Materials: As listed in the Confidential Instructions												
* 🚃	READ THESE INSTRUCTIONS FIRST												
	 Write your Centre number, candidate number and name on all the work you Give details of the practical session and laboratory where appropriate, in the Write in dark blue or black pen. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid. DO NOT WRITE IN ANY BARCODES. Answer all questions. Electronic calculators may be used. 												
	You may lose marks if you do not show your working or if you do not use appropriate units.												
	Use of a Data Booklet is unnecessary.	Session											
	Qualitative Analysis Notes are printed on pages 10 and 11. A copy of the Periodic Table is printed on page 12.												
	At the end of the examination, fasten all your work securely together.												
	The number of marks is given in brackets [] at the end of each question or part question.												
		For Examiner's Use											
		1											
		2											
		3											
		Total											

This document consists of **12** printed pages.



1 The concentration of hydrogen peroxide may be given in moldm⁻³ or as 'volume strength'. You will determine the concentration of hydrogen peroxide in moldm⁻³ and in 'volume strength' by a gas collection method.

Hydrogen peroxide decomposes to form water and oxygen. The reaction is much faster in the presence of a catalyst such as manganese(IV) oxide.

$$2H_2O_2(aq) \rightarrow 2H_2O(l) + O_2(g)$$


'Volume strength' is defined as the volume of oxygen in cm³ produced from the decomposition of 1.0 cm³ of hydrogen peroxide at room temperature and pressure. For example, 1.0 cm³ of '100 volume' hydrogen peroxide will produce 100 cm³ of oxygen.

FA 1 is a solution of hydrogen peroxide, H_2O_2 . **FA 2** is manganese(IV) oxide, MnO_2 .

(a) Method

Read the whole method before starting any practical work.

The diagram below may help you in setting up your apparatus.



- Fill the tub with water to a depth of about 5 cm.
- Fill the 250 cm³ measuring cylinder **completely** with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so that the open end is in the water just above the base of the tub.
- Rinse the 50 cm³ measuring cylinder with a little **FA1** then use it to transfer 150 cm³ of **FA1** into the reaction flask labelled **X**.
- Check that the bung fits tightly in the neck of flask **X**, clamp flask **X** and place the end of the delivery tube into the inverted 250 cm³ measuring cylinder.
- Remove the bung from the neck of the flask. Tip FA 2 into the hydrogen peroxide and replace the bung immediately. Remove the flask from the clamp and swirl it to mix the contents. Swirl the flask occasionally until no more gas is given off. Replace the flask in the clamp.
- Measure and record the final volume of gas in the measuring cylinder in the space below.

Keep FA 1 for use in Question 2.

Result

(b) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Use the information on page 2 to calculate the 'volume strength' of **FA 1**.

'volume strength' of **FA 1** =

(ii) Calculate the number of moles of oxygen collected in the measuring cylinder. [Assume 1 mole of gas occupies 24.0 dm³ under these conditions.]

moles of O_2 = mol

(iii) Using your answer to (ii) calculate the number of moles of hydrogen peroxide in the volume of FA 1 added to flask X.

moles of H_2O_2 = mol

(iv) Calculate the concentration of hydrogen peroxide, **FA 1**, in mol dm⁻³.

concentration of H_2O_2 , **FA 1 =** mol dm⁻³ [4] (c) (i) A source of error in this experiment is that some oxygen escapes before the bung can be inserted.

Suggest a change to the practical procedure given in **(a)** to reduce this source of error. You may draw a diagram as part of your answer.

(ii) The error in reading a 50 cm^3 measuring cylinder is $\pm 0.5 \text{ cm}^3$.

Calculate the maximum percentage error in the volume of hydrogen peroxide added to flask **X** in **(a)**.

maximum percentage error in volume of H_2O_2 = %

(iii) Explain why the presence of 20 cm³ of air in the 250 cm³ measuring cylinder before the start of the experiment would decrease the accuracy of the results obtained in (a).

[4]

(d) If you repeated the method described using half the mass of **FA 2**, what volume of gas would you expect to collect? Explain your answer.

[1] [Total: 11] 2 You will carry out a second experiment to determine the concentration of hydrogen peroxide, **FA 1**, in mol dm⁻³, by titration with acidified aqueous potassium manganate(VII). The equation for the reaction is given below.

 $2MnO_4^{-}(aq) + 5H_2O_2(aq) + 6H^{+}(aq) \rightarrow 2Mn^{2+}(aq) + 8H_2O(I) + 5O_2(g)$

FA 1 is a solution of hydrogen peroxide, H_2O_2 . **FA 3** is 0.0300 mol dm⁻³ potassium manganate(VII), KMnO₄.

FA 4 is dilute sulfuric acid.

(a) Method

- Fill the burette with **FA 3**.
- Pipette 25.0 cm³ of **FA 1** into a conical flask.
- Use the 25 cm³ measuring cylinder to add approximately 20 cm³ of **FA 4** to the conical flask.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is cm³.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of **FA 3** added in each accurate titration.

Ι	
II	
III	
IV	
V	
VI	
VII	

[7]

(b) From your accurate titration results, obtain a suitable value for the volume of **FA 3** to be used in your calculations. Show clearly how you obtained this value.

25.0 cm³ of **FA 1** required cm³ of **FA 3**. [1]

(c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of manganate(VII) ions present in the volume of FA 3 calculated in (b).

moles of MnO_4^- = mol

(ii) Calculate the number of moles of hydrogen peroxide present in 25.0 cm³ of **FA 1**.

moles of H_2O_2 = mol

(iii) Using your answer to (ii) calculate the concentration, in moldm⁻³, of hydrogen peroxide in **FA 1**.

concentration of H_2O_2 in **FA 1** = mol dm⁻³ [4]

[Total: 12]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs. No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) FA 5, FA 6 and FA 7 are solutions, some of which contain ions that are listed on pages 10 and 11.

		observations									
	test	FA 5	FA 6	FA 7							
(i)	To a 0.5 cm depth of solution in a boiling tube add aqueous sodium hydroxide, then										
	warm gently.										
	Allow to cool, add a piece of aluminium foil and warm again.										
(ii)	To a 1 cm depth of solution in a test-tube add two or three drops of acidified aqueous potassium manganate(VII). (Do not use FA 3 .)										
	If no reaction occurs, pour the mixture into a boiling tube and warm gently.										
(iii)	To a 1 cm depth of solution in a test-tube add a 2 cm depth of '10 volume' hydrogen peroxide and leave to stand. (Do not use FA 1 .)										
(iv)	To a 1 cm depth of solution in a test-tube add a 1 cm depth of dilute hydrochloric acid, then										
	add a 1 cm depth of aqueous barium chloride or aqueous barium nitrate.										

(b) (i) Identify as many ions present in **FA 5**, **FA 6** and **FA 7** as possible from your observations. If an ion cannot be identified from the tests, write 'unknown' in the space.

	cation(s)	anion(s)
FA 5		
FA 6		
FA 7		

(ii) Describe another test you could carry out to confirm the identity of a cation you have identified in (i). Record the reagent(s) and expected observation(s) in the space below.
 Do not carry out this test.

(iii) Write an ionic equation for the reaction that would occur in (ii). Include state symbols.

.....

[6]

[Total:17]

10

Qualitative Analysis Notes

1 Reactions of aqueous cations

ion	reaction with								
ion	NaOH(aq)	NH ₃ (aq)							
aluminium, A <i>l</i> ³⁺(aq)	white ppt. soluble in excess	white ppt. insoluble in excess							
ammonium, NH₄⁺(aq)	no ppt. ammonia produced on heating	_							
barium, Ba²⁺(aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.							
calcium, Ca²⁺(aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.							
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess							
copper(II), Cu²+(aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution							
iron(II), Fe²⁺(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess							
iron(III), Fe³⁺(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess							
magnesium, Mg²⁺(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess							
manganese(II), Mn²⁺(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess							
zinc, Zn²⁺(aq)	white ppt. soluble in excess	white ppt. soluble in excess							

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ^{2–}	CO ₂ liberated by dilute acids
chloride, C <i>l</i> ⁻(aq)	gives white ppt. with Ag ⁺ (aq) (soluble in $NH_3(aq)$)
bromide, Br⁻(aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in $NH_3(aq)$)
iodide, I⁻(aq)	gives yellow ppt. with Ag⁺(aq) (insoluble in NH₃(aq))
nitrate, NO₃⁻(aq)	NH_3 liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO₂⁻(aq)	NH ₃ liberated on heating with OH ⁻ (aq) and A <i>l</i> foil; NO liberated by dilute acids (colourless NO \rightarrow (pale) brown NO ₂ in air)
sulfate, SO₄²⁻(aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²-(aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO ₂)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H ₂	'pops' with a lighted splint
oxygen, O ₂	relights a glowing splint

1	2							GI	oup								
·			·									13	14	15	16	17	1
	Кеу																2 H heli 4
3	4			atomic number				1				5	6	7	8	9	1
Li	Be		ato	omic sym	bol							В	С	N	0	F	N
lithium 6.9	beryllium 9.0		rela	name ative atomic ma	255							boron 10.8	carbon 12.0	nitrogen 14.0	oxygen 16.0	fluorine 19.0	n 2
11	12											13	14	15	16	17	-
Na	Mg											Al	Si	Р	S	Cl	A
sodium 23.0	magnesium 24.3	3	4	5	6	7	8	9	10	11	12	aluminium 27.0	silicon 28.1	phosphorus 31.0	sulfur 32.1	chlorine 35.5	ai 3
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	
К	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	ł
potassium 39.1	calcium 40.1	scandium 45.0	titanium 47.9	vanadium 50.9	chromium 52.0	manganese 54.9	iron 55.8	cobalt 58.9	nickel 58.7	copper 63.5	zinc 65.4	gallium 69.7	germanium 72.6	arsenic 74.9	selenium 79.0	bromine 79.9	kr <u>i</u> 8
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	
Rb	Sr	Y	Zr	Nb	Мо	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I)
rubidium 85.5	strontium 87.6	yttrium 88.9	zirconium 91.2	niobium 92.9	molybdenum 95.9	technetium -	ruthenium 101.1	rhodium 102.9	palladium 106.4	silver 107.9	cadmium 112.4	indium 114.8	tin 118.7	antimony 121.8	tellurium 127.6	iodine 126.9	xe 13
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	
Cs	Ba	lanthanoids	Hf	Та	W	Re	Os	Ir	Pt	Au	Hg	Τl	Pb	Bi	Po	At	F
caesium 132.9	barium 137.3		hafnium 178.5	tantalum 180.9	tungsten 183.8	rhenium 186.2	osmium 190.2	iridium 192.2	platinum 195.1	gold 197.0	mercury 200.6	thallium 204.4	lead 207.2	bismuth 209.0	polonium —	astatine –	ra
87	88	89–103	104	105	106	107	108	109	110	111	112	20111	114	20010	116		
Fr	Ra	actinoids	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn		Fl		Lv		
francium –	radium —		rutherfordium	dubnium —	seaborgium	bohrium —	hassium —	meitnerium —	darmstadtium –	roentgenium	copernicium -		flerovium —		livermorium —		
		57	58	59	60	61	62	63	64	65	66	67	68	69	70	71]
lanthanoids		La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Ēr	Tm	Yb	Lu	
		lanthanum 138.9	cerium 140.1	praseodymium 140.9	neodymium 144.4	promethium —	samarium 150.4	europium 152.0	gadolinium 157.3	terbium 158.9	dysprosium 162.5	holmium 164.9	erbium 167.3	thulium 168.9	ytterbium 173.1	lutetium 175.0	
		89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	
ictinoids		Ac	Th	Pa	U uranium	Np	Pu	Am americium	Cm	Bk berkelium	Cf californium	Es einsteinium	Fm	Md	No nobelium	Lr lawrencium	

12