

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**MATHEMATICS**

**9709/53**

Paper 5 Mechanics 2 (**M2**)

**October/November 2017**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

Where a numerical value for the acceleration due to gravity is needed, use  $10 \text{ m s}^{-2}$ .

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [ ] at the end of each question or part question.

The total number of marks for this paper is 50.

This document consists of **13** printed pages and **3** blank pages.



1



A hollow cylinder with a rough inner surface has radius 0.5 m. A particle  $P$  of mass 0.4 kg is in contact with the inner surface of the cylinder. The particle and cylinder rotate together with angular speed  $6 \text{ rad s}^{-1}$  about the vertical axis of the cylinder, so that the particle moves in a horizontal circle (see diagram). Given that  $P$  is about to slip downwards, find the coefficient of friction between  $P$  and the surface of the cylinder. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



3



One end of a light inextensible string of length 0.4 m is attached to a fixed point  $A$  which is above a smooth horizontal surface. A particle  $P$  of mass 0.6 kg is attached to the other end of the string.  $P$  moves in a circle on the surface with constant speed  $v \text{ m s}^{-1}$ , with the string taut and making an angle of  $60^\circ$  with the horizontal (see diagram).

(i) Given that  $v = 0.5$ , calculate the magnitude of the force that the surface exerts on  $P$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the greatest possible value of  $v$  for which  $P$  remains in contact with the surface. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A particle  $P$  is projected with speed  $25 \text{ m s}^{-1}$  at an angle of  $30^\circ$  above the horizontal from a point  $O$  on horizontal ground. At time  $t$  s after projection the horizontal and vertically upwards displacements of  $P$  from  $O$  are  $x$  m and  $y$  m respectively.

(i) Express  $x$  and  $y$  in terms of  $t$  and hence show that the equation of the trajectory of  $P$  is

$$y = \frac{x}{\sqrt{3}} - \frac{4x^2}{375}. \quad [4]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- 5 One end of a light elastic string of natural length 0.8 m and modulus of elasticity 24 N is attached to a fixed point  $O$ . The other end of the string is attached to a particle  $P$  of mass 0.3 kg.  $P$  is projected vertically upwards with speed  $4 \text{ m s}^{-1}$  from a position 1.2 m vertically below  $O$ .

(i) Calculate the speed of the particle at the position where it is moving with zero acceleration. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





6 A solid object consists of a uniform hemisphere of radius 0.4 m attached to a uniform cylinder of radius 0.4 m so that the circumferences of their circular faces coincide. The hemisphere and cylinder each have weight 20 N. The centre of mass of the object lies at the centre  $O$  of their common circular face.

(i) Show that the height of the cylinder is 0.3 m. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A new object is made by cutting the cylinder in half and removing the half not attached to the hemisphere. The cut is perpendicular to the axis of symmetry, so the new object consists of a hemisphere and a cylinder half the height of the original cylinder.

(ii) Find the distance of the centre of mass of the new object from  $O$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The new object is placed with its hemispherical part on a rough horizontal surface. The new object is held in equilibrium by a force of magnitude  $P$  N acting along its axis of symmetry, which is inclined at  $30^\circ$  to the horizontal.

**(iii)** Find  $P$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 A particle  $P$  of mass  $0.2 \text{ kg}$  is released from rest at a point  $O$  on a rough plane inclined at  $60^\circ$  to the horizontal, and travels down a line of greatest slope. The coefficient of friction between  $P$  and the plane is  $0.3$ . A force of magnitude  $0.6x \text{ N}$  acts on  $P$  in the direction  $PO$ , where  $x \text{ m}$  is the displacement of  $P$  from  $O$ .

(i) Show that  $v \frac{dv}{dx} = 5\sqrt{3} - 1.5 - 3x$ , where  $v \text{ m s}^{-1}$  is the velocity of  $P$  at a displacement  $x \text{ m}$  from  $O$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the value of  $x$  for which  $P$  reaches its maximum velocity, and calculate this maximum velocity. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Calculate the magnitude of the acceleration of  $P$  immediately after it has first come to instantaneous rest. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.