

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs, tables or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions. A copy of the Periodic Table is printed on page 20.


At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part _ question.

| For Examin | |
|------------|--|
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| 8 | |
| 9 | |
| Total | |

This document consists of 20 printed pages.



(a) Fig. 1.1 is a side view of the thorax during breathing out and breathing in. The 1 and heart are not shown.



breathing out

breathing in



(i) Describe how each of the following have changed between breathing out and breathing in. the intercostal muscles the diaphragm [2] (ii) Explain how the changes you have described help to draw air into the lungs. [3] (b) As air is drawn into the lungs, it flows through the trachea and bronchi. These are lined with a tissue containing goblet cells and ciliated cells. Explain how this tissue helps to prevent infections in the lungs. [2]

| | | 3 | |
|-----|------|--|---------|
| (c) | Des | 3 scribe the effects of smoking on the goblet cells and cilia, | For |
| | (i) | the goblet cells and cilia, | mbridge |
| | | | |
| | | [2 | |
| | (ii) | the alveoli in the lungs. | |
| | | | |
| | | [2 | |

| | 4 |
|-------------------|---|
| emen | 4 nineteenth century, the Russian scientist Dimitri Mendeleev, arranged the its in order of the relative masses of their atoms. His work led to the modern Pen- hat we use today. Explain why atoms of different elements have different masses. |
| a) (i) | Explain why atoms of different elements have different masses. |
| | |
| | [1] |
| (ii) | Explain, in terms of electron configuration, why the element with proton number 36 is unreactive. |
| | [1] |
| (iii) | In the modern Periodic Table the elements with proton numbers 18 and 19 are not in order of their relative atomic masses. |
| | Suggest a reason for this. |
| | [1] |
| b) Ma | gnesium reacts with dilute hydrochloric acid according to the equation below. |
| | |
| | Mg + 2HC $l \longrightarrow$ MgC l_2 + H ₂ |
| | $\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$ |
| | student was asked to add 0.96 g of magnesium ribbon to 100 cm ³ of dilute |
| hyo | student was asked to add 0.96 g of magnesium ribbon to 100 cm ³ of dilute drochloric acid which had a concentration of 0.5 mol/dm ³ . |
| hyo | student was asked to add 0.96 g of magnesium ribbon to 100 cm ³ of dilute drochloric acid which had a concentration of 0.5 mol/dm ³ . Calculate the number of moles of magnesium in 0.96 g. Show your working. |
| hyo | student was asked to add 0.96 g of magnesium ribbon to 100 cm ³ of dilute drochloric acid which had a concentration of 0.5 mol/dm ³ . Calculate the number of moles of magnesium in 0.96 g. |
| һус (i) | student was asked to add 0.96 g of magnesium ribbon to 100 cm ³ of dilute drochloric acid which had a concentration of 0.5 mol/dm ³ . Calculate the number of moles of magnesium in 0.96 g. Show your working. [1] Calculate the number of moles of hydrochloric acid in 100 cm ³ of a solution which |
| һус (i) | student was asked to add 0.96 g of magnesium ribbon to 100 cm ³ of dilute drochloric acid which had a concentration of 0.5 mol/dm ³ . Calculate the number of moles of magnesium in 0.96 g. Show your working. [1] Calculate the number of moles of hydrochloric acid in 100 cm ³ of a solution which has a concentration of 0.5 mol/dm ³ . |

www.papaCambridge.com (iii) Use the balanced equation for this reaction and your results from (i) and predict whether there is enough acid to react with all of the magnesium.


..... [2]

(c) Fluorine is a halogen produced by electrolysis of an electrolyte containing fluoride ions, F⁻.

There were many attempts to produce fluorine during the nineteenth century and several scientists were seriously harmed when they succeeded in making fluorine. They attempted to collect fluorine in containers made of gold or platinum and they kept the containers at a very low temperature.


(i) State and explain at which electrode, cathode or anode, fluorine is produced during electrolysis.

[2] (ii) Use your knowledge of the halogen group to suggest why fluorine caused harm to scientists who first produced it. (iii) Suggest why the scientists attempting to produce fluorine used gold or platinum containers at a very low temperature. [2]



www.papaCambridge.com (b) A pedestrian steps into the path of the moving car. Fig. 3.2 shows a graph of h speed of the car changes from the moment when the driver sees the pedestrian the car stops.

7





(i) After 1.2 s the car slows down.

Calculate the deceleration of the car.

State the formula that you use and show your working.

formula used


working

- [2]
- (ii) Calculate the total distance travelled by the car between the driver seeing the pedestrian and the car stopping.

Show your working.

[3]

An experiment was carried out into the effect of different doses of X-rays on the spen 4 produced by male fruit flies. Fig. 4.1 shows the results.



[2]

www.papaCambridge.com 9 (c) Fruit flies have four pairs of chromosomes in their cells. Some of the mutations in the experiment above involved the loss of one chromosome If a fruit fly sperm that had lost one chromosome fertilised a normal egg, how many chromosomes would there be in the zygote? [1] . (d) Explain why a mutation that occurs in a gamete-forming cell is more likely to be harmful than one that occurs elsewhere in a fruit fly's body. _____ [2] (e) Insects can be serious pests, for example by carrying disease or eating crops. Pesticides can be used to kill them, but many people are concerned about the harm that pesticides do and are trying other methods of controlling insect populations. One new method that is being tested is to expose a large number of male insects of a harmful species to X-rays and then release them into the wild. (i) Explain why people are concerned about the use of pesticides. [2] (ii) Suggest how the new method might reduce the population of the harmful insects. [2]

www.papaCambridge.com 10 (a) Glucose and starch are carbohydrates. 5 (i) The chemical formula of glucose is $C_6H_{12}O_6$. State the total number of atoms which are combined in one molecule of glucose. (ii) Explain why it is not possible to write a simple chemical formula for starch. [2] (b) Fig. 5.1 shows an experiment which was set up to investigate the action of a partially permeable membrane. A tube made from a partially permeable membrane was filled with iodine solution and placed into a beaker containing a mixture of glucose, starch and water. mixture of glucose, iodine solution starch and water tube made from partially permeable membrane Fig. 5.1 (i) Explain the following observations which were made some time later.

The solution **inside** the tube gave a positive result with Benedict's solution.

The solution **outside** the tube became blue-black in colour. [4] (ii) Predict and explain whether the solution inside the tube became blue-b colour.

www.papacambridge.com

(c) Plastics are materials made mainly from polymer molecules. Fig. 5.2 shows part of a polymer molecule. Molecules of this polymer are formed by addition polymerisation of an unsaturated monomer.




Fig. 5.2

(i) Draw the displayed formula of one of the monomer molecules which have joined to form this polymer.

[2]

(ii) Two different plastics, A and B, were heated. Plastic A melted easily but plastic B did not melt even when heated to a very high temperature.

Explain these observations. You may draw some simple diagrams to help your answer.

..... [3]

6 Fig. 6.1 shows a circuit containing four ammeters, A_1 , A_2 , A_3 and A_4 .




Fig. 6.1

Table 6.1 shows the readings on each ammeter.

| ammeter | reading on ammeter / amps |
|-----------------------|------------------------------|
| A ₁ | |
| A ₂ | 0.2 |
| A ₃ | 0.3 |
| A ₄ | 0.5 |

(a) What is the reading on ammeter A_1 ?

[1]

(b) Calculate the combined resistance of the two resistors in the circuit in Fig. 6.1.

State the formula that you use and show your working.

formula used

working

[3]

www.papacambridge.com

- (c) Fig. 6.2 shows a magnet and coil of wire connected to a sensitive ammeter.

sensitive ammeter

Fig. 6.2

(i) When the magnet is moved into the coil, the needle on the ammeter shows a deflection to the left.


Explain why a reading on the ammeter is produced.


[2]


(ii) Explain how this effect is used in a dynamo to produce an output voltage. You may use a diagram to help with your answer.

| [4] |
|-----|
| |

www.papaCambridge.com







Describe and explain how water from the soil is transported up to the leaves at the top of the tree.

| [3] |
|-----|

www.papacambridge.com

- 8 In many countries supplies of clean water for drinking are obtained from river water.
 - (a) State two processes that are used to convert river water into water which is safe humans to drink.
- www.papaCambridge.com 1. 2. [2]
 - (b) A sample of safe drinking water still contained dissolved calcium sulphate, CaSO₄. which helped to make the water hard.
 - (i) State the formula of the particle present in this water which causes hardness.

[1]

(ii) A student carried out an experiment to find out if boiling would remove the hardness from this sample of water.

The results of his experiment are shown in Table 8.1.

| Table 8.1 | |
|-----------|--|
|-----------|--|

| water sample | volume of water tested / cm ³ | volume of soap solution needed for lather / cm ³ |
|----------------------------------|---|---|
| distilled water | 25.0 | 0.2 |
| hard water control (unboiled) | 25.0 | 8.0 |
| hard water boiled for 5 minutes | 25.0 | 3.0 |
| hard water boiled for 10 minutes | 25.0 | 3.0 |

What conclusions could the student draw from these results?

[2]

- (c) Some types of salt used to flavour food are mixtures of sodium chloride and poly chloride. Sodium chloride and potassium chloride are both ionic compounds.
- www.papaCambridge.com (i) Potassium chloride can be formed by reacting potassium directly with chlorine. Fig. 8.1 shows the electron arrangements in a potassium atom and a chlorine atom.




Fig. 8.1

In the space below, draw diagrams similar to those in Fig. 8.1 which show the electron arrangements of the two particles when combined in potassium chloride.

[2]

(ii) Explain briefly why potassium chloride is a solid with a high melting point at room temperature.

..... [2]

| | the second second | |
|---------|---|------------|
| | 18 | 1 |
| A polic | e car uses a siren and a blue light to alert people. | Cal |
| (a) (i) | 18 e car uses a siren and a blue light to alert people. Explain why sound needs a medium, such as air, to travel through. | |
| | | |
| (ii) | How will the sound of the siren change if the amplitude of the sound waves emities increased? | [2] ted |
| | | [1] |
| (iii) | Suggest a suitable frequency for the sound emitted by the siren to alert people. | |
| | | [1] |
| | e police communicate using radio waves. Both blue light and radio waves are part electromagnetic spectrum. | t of |
| (i) | State one property which all electromagnetic waves have in common. | |
| (ii) | State one difference between blue light waves and radio waves. | [1] |
| (ii) | | [1] |
| (iii) | The radio waves used have a frequency of 10 000 000 Hz and a wavelength 30 m. | |
| | Calculate the speed of these waves. | |
| | State the formula that you use and show your working. | |
| | formula used | |
| | working | |
| | | |
| | | |

- (c) As the police car drives along the temperature of the air in the tyres increases.
 - (i) Use the ideas of the kinetic theory to explain why this will result in an increase tyre pressure.

www.papaCambridge.com [2]

(ii) The original temperature of the air in the tyres was 10 °C and the final temperature was 30°C.

Calculate the final pressure of the air in the tyres if the original pressure was $200\ 000\ \text{N/m}^2$.

State the formula that you use and show your working.

formula used

working

[3]

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of

| Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity. |
|--|
|--|

| | 1 | | | | | | | Gr | oup | | | | | 1 | | | | _ |
|----------------------------------|------------------------------------|--|--|---|---------------------------------|-----------------------------------|-------------------------------------|------------------------------------|-------------------------------------|--------------------------------------|-----------------------------------|-------------------------------------|-----------------------------|---|--|-------------------------------------|-------------------------------|-------|
| | II | | | | | | | | | | | | IV | V | VI | VII | 0 | |
| | | | | | | | 1 H Hydrogen 1 | | | | | | | | | | 4 He Helium | |
| 7 Li Lithium | 9 Be Beryllium | | | | | | | _ | | | | 11 B Boron 5 | 12 C Carbon 6 | 14 N Nitrogen | 16 O Oxygen 8 | 19 F Fluorine 9 | 20 Ne Neon 10 | |
| 23 Na Sodium | 24 Mg Magnesium 12 | | | | | | | | | | | 27 A 1 Aluminium 13 | 28 Si Silicon 14 | 31 P Phosphorus 15 | 32 S Sulphur 16 | 35.5 C1 Chlorine 17 | 40 Ar ^{Argon} | |
| 39 K Potassium 9 | 40 Ca Calcium 20 | 45 Sc Scandium 21 | 48 Ti Titanium 22 | 51 V Vanadium 23 | 52 Cr Chromium 24 | 55 Mn Manganese 25 | 56 Fe Iron 26 | 59 Co Cobalt 27 | 59 Ni _{Nickel} 28 | 64 Cu Copper 29 | 65 Zn Zinc 30 | 70 Ga Gallium 31 | 73 Ge Germanium 32 | 75 As Arsenic 33 | 79 Se Selenium 34 | 80 Br Bromine 35 | 84 Kr Krypton 36 | 20 |
| 85 Rb Rubidium 7 | 88 Sr Strontium 38 | 89 Y Yttrium 39 | 91 Zr Zirconium 40 | 93 Nb Niobium 41 | 96 Mo Molybdenum 42 | Tc Technetium 43 | 101 Ru Ruthenium 44 | 103 Rh Rhodium 45 | 106 Pd Palladium 46 | 108 Ag Silver 47 | 112 Cd Cadmium 48 | 115 In Indium 49 | 119 Sn 50 | 122 Sb Antimony 51 | 128 Te Tellurium 52 | 127 I Iodine 53 | 131 Xe Xenon 54 | 0 |
| 133 Cs Caesium 5 | 137 Ba Barium 56 | 139 La Lanthanum 57 * | 178 Hf ^{Hafnium} 72 | 181 Ta Tantalum 73 | 184 W Tungsten 74 | 186 Re Rhenium 75 | 190 Os Osmium 76 | 192 Ir Iridium 77 | 195 Pt Platinum 78 | 197 Au Gold 79 | 201 Hg Mercury 80 | 204 T 1 Thallium 81 | 207 Pb Lead 82 | 209 Bi Bismuth 83 | Polonium 84 | At Astatine 85 | Rn Radon 86 | |
| Fr Francium 7 | 226 Ra Radium 88 | 227 Ac Actinium 89 † | | | | | | | | | | | | | | | | |
| | anthanoi Actinoid | series | | 140 Ce ^{Cerium} 58 | 141 Pr Praseodymium 59 | 144 Nd Neodymium 60 | Pm Promethium 61 | 150 Sm Samarium 62 | 152 Eu Europium 63 | 157 Gd Gadolinium 64 | 159 Tb Terbium 65 | 162 Dy Dysprosium 66 | 165 Ho Holmium 67 | 167 Er ^{Erbium} 68 | 169 Tm ^{Thulium} 69 | 173 Yb Ytterbium 70 | 175 Lu Lutetium 71 | |
| ey Þ | x x | = relative atom = atomic symb = proton (atom | bol | 232 Th Thorium 90 | Pa Protactinium 91 | 238 U Uranium 92 | Np Neptunium 93 | Pu Plutonium 94 | Am Americium 95 | Cm Curium 96 | Bk Berkelium 97 | Cf Californium 98 | Es Einsteinium 99 | Fm Fermium 100 | Md Mendelevium 101 | No Nobelium 102 | Lr Lawrencium 103 | |
| | | | | The v | volume of | one mole | of any ga | as is 24 d | m ³ at roo | m temper | ature and | l pressure | (r.t.p.). | | | | 71 Lr Lawrencium 103 | Jedeo |