



CO-ORDINATED SCIENCES

0654/41

Paper 4 Theory (Extended)

May/June 2017

MARK SCHEME

Maximum Mark: 120

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

© IGCSE is a registered trademark.

This document consists of **11** printed pages.

Question	Answer	Marks
1(a)(i)	Y umbilical cord ; Z amniotic fluid ;	2
1(a)(ii)	protects (the fetus) from mechanical shock ; protects (the fetus) from drying out ; protects (the fetus) from temperature fluctuations ;	max 1
1(a)(iii)	carries oxygen / glucose / nutrients, to the fetus ; carries, urea / toxins / carbon dioxide / waste products, away from fetus ;	2
1(b)(i)	accept an 'X' placed anywhere immediately above or on the top of the cervix ;	1
1(b)(ii)	bleeding / haemorrhaging ; damage to placenta ; blocks passage of baby / AW ;	max 1

Question	Answer	Marks
2(a)(i)	temperature change = $31\text{ }^{\circ}\text{C}$; $E/m\Delta\theta$ / $156\ 000 / 1.2 \times 31$; = $4190 / 4194$ (J / Kg $^{\circ}\text{C}$) ;	3
2(a)(ii)	efficiency = useful energy out / energy in $\times 100$ / $2600 / 3000 \times 100$; = $87(\%)$;	2
2(b)	latent heat (of vaporisation) required ; as energy to break bonds / to overcome attractive forces ; between molecules / intermolecular bonds ; to increase <u>potential</u> energy of the molecules ;	2

Question	Answer	Marks
3(a)(i)	produced in car engines / by lightning ; contributes to acid rain / acidifies lakes / reference to damage to, plants / aquatic organisms / reference to damage to (animal) respiratory systems / damage to buildings / AVP ;	2
3(a)(ii)	Haber ;	1
3(a)(iii)	$\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightarrow \text{CO}(\text{g}) + 3 \text{H}_2(\text{g})$ symbols and state symbols ; balanced ;	2
3(b)(i)	6 shared electrons ; remaining lone pair ;	2
3(b)(ii)	multiple bonding / 6 / 3 pairs, bonding electrons / triple bond ; bond between the atoms is very strong / difficult to break / (relatively) large amount of energy required (to break bond) ;	2
3(c)(i)	M_r of hydrazine = $(14 \times 2) + (1 \times 4)$;	1
3(c)(ii)	moles of hydrazine = $192 \div 32 = 6$; so moles of ammonia = $4 \times 2 / 4 \times 6 \div 3 = 8$; volume of ammonia = $8 \times 24 = 192 \text{ (dm}^3\text{)}$;	3

Question	Answer	Marks
4(a)(i)	(pH) 9 ;	1
4(a)(ii)	enzyme, is denatured / changes shape ;	1
4(a)(iii)	temperature / substrate concentration ;	1
4(b)	breakdown of large molecules into small molecules ; from insoluble to soluble ; using, mechanical / chemical, processes / means ;	3
4(c)	(enzyme) A ; (enzyme A) works at low pH / in acidic conditions / optimum pH is 1.9 ;	2

Question	Answer	Marks
5(a)	$6\text{CO}_2 + 6\text{H}_2\text{O} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2$ symbols ; balancing ;	2
5(b)	stunted weak stem ;	1
5(c)(i)	transpiration ; water loss / diffusion of water vapour / evaporation from, leaf / stomata ; (water and ions) drawn up xylem ; down water potential gradient ; ref to cohesion of water molecules ;	max 3
5(c)(ii)	less transpiration / diffusion of water vapour / water loss / evaporation ; smaller water potential gradient ; slower movement of, water / ions ;	max 2
5(d)	eutrophication ; algal bloom causes lack of light ; lack of light causes death of plants ; death of plants causes increase in bacteria ; increase in bacteria / bacteria respiration, reduces oxygen concentration reduced oxygen kills fish ;	max 3

Question	Answer	Marks
6(a)(i)	<u>fractional distillation</u> ;	1
6(a)(ii)	(average) size / surface area of molecules increases ; so intermolecular forces / forces between molecules increase ; so greater (thermal) energy / higher temperature required to separate molecules ;	3
6(a)(iii)	pure / single substances have discrete boiling point / owtte ; liquid mixture has a range of boiling point ;	2
6(b)(i)	<pre> H H H H H — C — C — C — C — H H H H H </pre> 4 carbon atoms and 10 hydrogen atoms ; all else correct ;	2
6(b)(ii)	flammable / produce CO ₂ / H ₂ O / CO when burnt ;	1

Question	Answer	Marks
7(a)(i)	4 (m / s) ;	1
7(a)(ii)	area under graph / working ; 20 + 20 + 50 = 90 (m) ;	2
7(a)(iii)	working ; e.g. correct substitution into formula such as 4 / 10 ;	1
7(a)(iv)	force = mass \times acceleration / 950×0.4 ; 380 (N) ;	2
7(b)(i)	move faster ;	1
7(b)(ii)	more frequent collisions / collide at greater speed, with tyre wall ; more force exerted on tyre walls ;	2
7(c)(i)	current in low voltage circuit creates magnetic field (around solenoid) ; soft iron attracted (to magnet / solenoid) ; contacts in high voltage circuit close ;	3
7(c)(ii)	so that humans, are not exposed to the high voltage circuit / operate low voltage switching circuit / owtte ;	1

Question	Answer	Marks					
8(a)(i)	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>(D)</td> <td>E</td> <td>(A)</td> <td>C</td> <td>B</td> </tr> </table> ;	(D)	E	(A)	C	B	1
(D)	E	(A)	C	B			
8(a)(ii)	brain is closer / spinal cord is further away ;	1					
8(b)(i)	radial muscles contract ; pupil widens ; let more light into the, eye / retina ;	max 2					
8(b)(ii)	no conscious thought / automatic / immediate / rapid ;	1					
8(c)	bigger eyes / wider pupils ;	1					

Question	Answer	Marks
9(a)(i)	12 protons ; 14 neutrons ;	2
9(a)(ii)	2,8,2 ;	1
9(b)(i)	hydrogen ;	1
9(b)(ii)	(concentration of) acid decreases ; (concentration of) magnesium chloride increases ; (mass of) magnesium decreases ;	max 2
9(b)(iii)	K.E. higher at B than at A AND K.E. at B and C the same ;	1
9(c)(i)	reaction releases thermal energy / temperature of mixture increases / exothermic / temperature affects rate ; water (seeks to) keep temperature constant ;	2
9(c)(ii)	reaction rate increases ; increased collision frequency ;	2

Question	Answer	Marks
10(a)	suitable temperature / warmth AND water / moisture ;	1
10(b)	glucose ;	1
10(c)	red liquid would move, further / more quickly (to the left) ; increased respiration ; increased oxygen used ;	3
10(d)	no movement of red liquid ; enzymes denatured ; no respiration / no oxygen used ;	3

Question	Answer	Marks
11(a)	use Geiger counter etc. ; test for absorption by shield of lead / thick aluminium ; γ -rays are more penetrating than α or β / α and β will not penetrate lead ; OR measure deflection by magnetic / electric field ; γ -rays not deflected / α and β deflected ;	3
11(b)	${}_{92}^{235}\text{U}$; ${}_{2}^{4}\text{He}$ OR ${}_{2}^{4}\alpha$;	2
11(c)	correct working ; 28 (Ω) ;	2
11(d)(i)	approx sin wave ; constant amplitude ;	2
11(d)(ii)	stronger magnet / spin coil faster / greater number of turns / increased coil area ;	1
11(e)(i)	$\lambda = v / f / 340 / 490$; $= 0.69$ (m) ;	2
11(e)(ii)	compression correctly labelled ;	1
11(e)(iii)	decreases / closer together ;	1

Question	Answer	Marks
12(a)(i)	sodium atoms lose one electron / change from 2,8,1 to 2,8 ; chlorine atoms gain one electron / change from 2,8,7 to 2,8,8 ;	2
12(a)(ii)	alternating sodium and chloride ions in two directions ;	1
12(b)(i)	(aqueous NaCl) hydrogen chlorine ; (molten NaCl) sodium chlorine ;	2
12(b)(ii)	mobile ions carry charge / produce current / allow electricity to flow ; ions are not mobile / fixed in a solid ;	2

Question	Answer	Marks
13(a)(i)	light travels faster than sound ;	1
13(a)(ii)	region where a charge experiences a force ;	1
13(a)(iii)	current = charge / time / 1.21 / 0.00011 ; = 11000 (A) ;	2
13(b)(i)	middle ray passes through without deviation AND bottom ray passes out parallel to principal axis AND all 3 rays pass through a point ;	1
13(b)(ii)	inverted arrow drawn from principal axis to intersection of three rays ;	1