

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/53

Paper 5 Practical Test

October/November 2021

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use				
1				
2				
3				
Total				

This document has 12 pages. Any blank pages are indicated.

1 You are going to investigate the temperature change when zinc reacts with two different aqueous solutions of copper(II) sulfate, solution **Q** and solution **R**.

Read all of the instructions carefully before starting the experiments.

Instructions

You are going to do two experiments.

(a) Experiment 1

- Place a polystyrene cup into a 250 cm³ beaker for support.
- Use a measuring cylinder to pour 25 cm³ of solution **Q** into the polystyrene cup.
- Use a thermometer to measure the initial temperature of solution **Q**. Record this in the space above the table.
- Add 3g of zinc powder to the polystyrene cup. At the same time start a stop-watch.
- Using the thermometer, continually stir the mixture in the polystyrene cup. Record the temperature every 30 seconds for 240 seconds. Record the temperatures in the table.

initial temperature =°C

time/s	30	60	90	120	150	180	210	240
temperature/°C								
temperature change/°C								

• Complete the table by calculating the temperature changes from the initial temperature using the equation:

temperatu	re change	= temperature	e – initial te	mperature

[3]

(b) Experiment 2


- Empty the polystyrene cup and rinse with distilled water.
- Repeat Experiment 1 using solution R in place of solution Q.
- Complete the table by calculating the temperature changes from the initial temperature.

initial temperature =°C

time/s	30	60	90	120	150	180	210	240
temperature/°C								
temperature change/°C								

[3]

(c) Complete a suitable scale on the *y*-axis and plot your results from Experiment 1 and Experiment 2 on the grid. Draw two curves of best fit. Both curves must start at (0,0). Label your curves.



[5]

(d) From your graph, deduce the temperature change at 110 seconds in Experiment 1.

Show clearly **on the grid** how you worked out your answer.

													0	\sim	[2	1
 	 	_	 	 	_	_		 	_	_	 				12	

(e) Predict the temperature of the solution in Experiment 2 after 5 hours. Explain your answer.

(i)	Suggest why the experiments were done in a polystyrene cup rather than a glass beak	er.
		[1]
(ii)	Describe how the results would be different if a glass beaker is used in place of t polystyrene cup.	:he
		[1]
		of
cha	inge	
ехр	lanation	
		 [2]
	(iii) Sugathe	(ii) Describe how the results would be different if a glass beaker is used in place of t polystyrene cup. Suggest one change that could be made to the apparatus that would improve the accuracy the results. Explain why this change would improve the accuracy of the results. change

[Total: 19]

2 You are provided with solid **S** and solid **T**.

Do the following tests on the substances, recording all of your observations at each stage.

(a) To solid **S** in the boiling tube, add about 10 cm³ of dilute hydrochloric acid.

tests on solid S

	rest arry gas produced.
	Keep the product for (b).
	Record your observations.
	[2]
(b)	The solution formed in (a) is solution U . Decant about 1 cm depth of solution U into a test-tube.
	To solution ${\bf U}$ add aqueous sodium hydroxide dropwise and then in excess. Record your observations.
	[2]
(c)	Identify solid S .

tests on solid T

(d)		solid ${\bf T}$ in the boiling tube, add about $10{\rm cm}^3$ of distilled water. Place a stopper in the boiling e and shake the tube to dissolve solid ${\bf T}$ and form solution ${\bf T}$.
	Div	ide solution T into four approximately equal portions in four test-tubes.
	(i)	To the first portion of solution \mathbf{T} , add aqueous sodium hydroxide dropwise and then in excess. Record your observations.
		[2]
	(ii)	Pour the second portion of solution T into the test-tube containing sodium carbonate. Record your observations.
	(iii)	To the third portion of solution T , add a 2cm length of magnesium ribbon. Record your observations.
		[2]
	(iv)	To the fourth portion of solution T , add 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate. Leave the mixture to stand for 5 minutes. Record your observations.
		[1]
(e)	lde	ntify solid T .

[Total: 15]

3	Catalysts are substances which increase the rate of a reaction but are unchanged at the end of the
	reaction.

Aqueous hydrogen peroxide decomposes slowly to form water and oxygen.

$$2H_2O_2(aq) \rightarrow 2H_2O(I) + O_2(g)$$

Copper(II) oxide is an insoluble solid.

Plan an investigation to find out if copper(II) oxide is a catalyst for the decomposition of hydrogen peroxide. You must include how your results will tell you if copper(II) oxide is a catalyst. You have access to copper(II) oxide, aqueous hydrogen peroxide and all normal laboratory apparatus.

BLANK PAGE

BLANK PAGE

BLANK PAGE

Notes for use in qualitative analysis Tests for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide (Br ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide, then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify, then add aqueous barium nitrate	white ppt.
sulfite (SO ₃ ²⁻)	add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide	sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless

Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+)	white ppt., soluble in excess, giving a colourless solution	white ppt., insoluble in excess
ammonium (NH ₄ ⁺)	ammonia produced on warming	-
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
chromium(III) (Cr ³⁺)	green ppt., soluble in excess	grey-green ppt., insoluble in excess
copper(II) (Cu ²⁺)	light blue ppt., insoluble in excess light blue ppt., soluble in excess, giving a dark blue solution	
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution

Tests for gases

gas	test and test result	
ammonia (NH ₃)	turns damp red litmus paper blue	
carbon dioxide (CO ₂)	turns limewater milky	
chlorine (Cl ₂)	bleaches damp litmus paper	
hydrogen (H ₂)	'pops' with a lighted splint	
oxygen (O ₂)	relights a glowing splint	
sulfur dioxide (SO ₂)	turns acidified aqueous potassium manganate(VII) from purple to colourless	

Flame tests for metal ions

metal ion	flame colour
lithium (Li ⁺)	red
sodium (Na ⁺)	yellow
potassium (K⁺)	lilac
copper(II) (Cu ²⁺)	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.