

# **Cambridge IGCSE**<sup>™</sup>

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/51

Paper 5 Practical Test

May/June 2022

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

#### **INSTRUCTIONS**

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

#### **INFORMATION**

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [ ].
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use		
1		
2		
3		
Total		


This document has 12 pages. Any blank pages are indicated.

1 You are going to investigate the rate at which hydrogen gas is made when magnesium reacts with two different solutions of dilute hydrochloric acid, **C** and **D**, with different concentrations. The dilute hydrochloric acid is in excess in both experiments.

## Read all of the instructions carefully before starting the experiments.

#### Instructions

You are going to do two experiments using the apparatus shown.



#### (a) Experiment 1

- Use a measuring cylinder to pour 50 cm³ of dilute hydrochloric acid **C** into a conical flask.
- Use a thermometer to measure the initial temperature of the dilute hydrochloric acid. Record the initial temperature in the space provided.
- Set the apparatus up as shown in the diagram, ensuring the inverted measuring cylinder is full of water.
- Remove the bung from the conical flask, leaving the delivery tube in the measuring cylinder.
- Add a coiled 5 cm length of magnesium ribbon to the conical flask, immediately replace the bung and start the timer.
- Measure the volume of gas collected in the inverted measuring cylinder every 20 seconds for 160 seconds. Record the volume of gas collected in the table.
- Use the thermometer to measure the final temperature of the dilute hydrochloric acid in the conical flask. Record the final temperature in the space provided.

initial temperature	°C final temperature							°C		
time/s	20	40	60	80	100	120	140	160		
volume of gas collected/cm <sup>3</sup>										

[2]

### (b) Experiment 2


- Empty and rinse the conical flask with distilled water.
- Repeat Experiment 1 using 50 cm³ of dilute hydrochloric acid **D** instead of dilute hydrochloric acid **C**.

initial temperature ......°C final temperature ......°C

time/s	20	40	60	80	100	120	140	160
volume of gas collected/cm³								

**(c)** Complete a suitable scale on the *y*-axis and plot your results from Experiments 1 and 2 on the grid.

Draw **two** smooth line graphs. The lines must pass through (0,0). Clearly label your lines.



(d)		om your graph, periment 2.	deduce the	volume o	of gas	that was	collected	after 50	seconds in
	Sho	ow clearly <b>on the 9</b>	grid how you	worked o	ut your	answer.			
						volume	of gas = .		[3]
(e)	hyd	olain what can be o drochloric acid <b>D</b> .					·		
									[2]
(f)	(i)	State what happe				·			•
	(ii)	State what effect reaction has finis	this temperat						
									[1]
	(iii)	Describe a chang change of the aci			ne appa	ratus or re	agents to re	educe the	temperature
									[1]
(g)		ggest why it is imp magnesium ribbo	n.						
(h)		ate the advantage ery 20 seconds.	of measuring	the volur	me of g	as collecte	ed every 1	0 seconds	s rather than
									[1]
									[Total: 20]

You are provided with two substances, solid E and solution F.
Do the following tests, recording all of your observations at each stage.

#### tests on solid E

Add about 15 cm³ of distilled water to the boiling tube containing solid **E**. Replace the stopper in the boiling tube and shake the boiling tube to dissolve solid **E** and form solution **E**. Divide solution **E** into three approximately equal portions in one boiling tube and two test-tubes.

(a) To the first portion of solution E in the boiling tube, add aqueous sodium hydroxide dropwise

	until it is in excess.	
	Keep the mixture formed for (b).	
	Record your observations.	
		[2]
(b)	Gently warm the mixture formed in <b>(a)</b> . Test any gas produced and identify the gas. Record your observations.	
	identity of gas	 [2]
(c)	To the second portion of solution <b>E</b> add about 1 cm depth of dilute nitric acid followed by a tdrops of aqueous silver nitrate.  Record your observations.	few
		[1]
(d)	To the third portion of solution <b>E</b> add about 1 cm depth of dilute nitric acid followed by a drops of aqueous barium nitrate.  Record your observations.	few
		[1]
(e)	Identify the <b>three</b> ions in solution <b>E</b> .	

# tests on solution F

	Divide solution F	into two	approximately	equal /	portions	in two	test-tubes.
--	-------------------	----------	---------------	---------	----------	--------	-------------

(f)	Test the pH of the first portion of solution <b>F</b> .
	pH = [1]
(g)	Add the second portion of solution <b>F</b> to the boiling tube containing solid sodium carbonate. Test any gas produced. Record your observations.
	[3]
(h)	Identify the positive ion in solution <b>F</b> .
	[1]
	[Total: 14]

A sample of muddy river water contains water, dissolved solids and insoluble solid mud.	
Plan an investigation to find the concentration of dissolved solids, in g/dm³, in the river water.	
In your answer state how you will work out the concentration of the <b>dissolved</b> solids in g/dm	3.
You are provided with a small sample (less than $1dm^3$ ) of muddy river water and common labora apparatus. ( $1dm^3=1000cm^3$ )	atory
	. [6]

# **BLANK PAGE**

# **BLANK PAGE**

# **BLANK PAGE**

# Notes for use in qualitative analysis Tests for anions

anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide (Br <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO <sub>3</sub> <sup>-</sup> ) [in solution]	add aqueous sodium hydroxide, then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-</sup> ) [in solution]	acidify, then add aqueous barium nitrate	white ppt.
sulfite (SO <sub>3</sub> <sup>2-</sup> )	add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide	sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless

# Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+)	white ppt., soluble in excess, giving a colourless solution	white ppt., insoluble in excess
ammonium (NH <sub>4</sub> <sup>+</sup> )	ammonia produced on warming	-
calcium (Ca <sup>2+</sup> )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
chromium(III) (Cr <sup>3+</sup> )	green ppt., soluble in excess	grey-green ppt., insoluble in excess
copper(II) (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess, giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution

#### **Tests for gases**

gas	test and test result
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint
sulfur dioxide (SO <sub>2</sub> )	turns acidified aqueous potassium manganate(VII) from purple to colourless

#### Flame tests for metal ions

metal ion	flame colour
lithium (Li <sup>+</sup> )	red
sodium (Na <sup>+</sup> )	yellow
potassium (K⁺)	lilac
copper(II) (Cu <sup>2+</sup> )	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.